arXiv:1609.04924v1 [math.NT] 16 Sep 2016

On harmonic numbers and nonlinear Euler sums

Ce Xu* Yulin Cai

School of Mathematical Sciences, Xiamen University

Xiamen 361005, P.R. China

Abstract In this paper we are interested in Euler-type sums with products of harmonic numbers, Stirling numbers and Bell numbers. We discuss the analytic representations of Euler-type sums through polylogarithm function and Riemann zeta function, and some numerical values of Euler sums are given. Furthermore, we give explicit formula for several classes of Euler-related sums in terms of zeta values and harmonic numbers, and several examples are given. The given representations are new.

Keywords Harmonic number; Stirling number; Euler sum; Riemann zeta function.

AMS Subject Classifications (2010): 11M06; 11M32; 33B15

Contents

1 Introduction **1**

2 Series involving Stirling numbers, harmonic numbers and Bell numbers **7**

3 Some identities On Euler Sums **24**

4 Triple Euler sums **34**

5 Some evaluation of Euler-type sums **40**

1 Introduction

Throughout this article we will use the following definitions and notations. In this paper, the generalized nth-harmonic numbers of order r are given by

$$\zeta_n(r) = \sum_{j=1}^{n} \frac{1}{j^r}, H_n = \zeta_n(1) = \sum_{j=1}^{n} \frac{1}{j}, \ 1 \leq r \in \mathbb{Z}, \tag{1.1}$$

where the empty sum $\zeta_0(r)$ is conventionally understood to be zero. These are given in terms of polygamma functions $\psi(r)$ as

$$\zeta_n(r) = \frac{(-1)^{r-1}}{(r-1)!}\left(\psi^{(r-1)}(n+1) - \psi^{(r-1)}(1)\right), H_n = \int_0^1 \frac{1-t^n}{1-t}dt = \psi(n+1) + \gamma,$$

*Corresponding author. Email: 15959259051@163.com (C. XU)

where $\psi(z)=\dfrac{d}{dz}(\ln\Gamma(z))=\dfrac{\Gamma'(z)}{\Gamma(z)}$ is digamma function and $\psi^{(n)}(z)=(-1)^{n+1}n!\sum_{k=0}^{\infty}1/(z+k)^{n+1}$, $\Gamma(z)=\int_0^{\infty}e^{-t}t^{z-1}dt,\ \Re(z)>0$ is gamma function, $\psi^{(r-1)}(1)=(-1)^r(r-1)!\zeta(r)$, γ denotes the Euler-Mascheroni constant defined by

$$\gamma=\lim_{n\to\infty}\left(\sum_{k=1}^{n}\frac{1}{k}-\ln n\right)=-\psi(1)\approx 0.57721566490153286060606512....$$

More generally we can define the generalized αth $(0<\alpha\in\mathbb{R})$ harmonic numbers of order r by

$$\zeta_{\alpha}(r)=\frac{(-1)^{r-1}}{(r-1)!}\left(\psi^{(r-1)}(\alpha+1)-\psi^{(r-1)}(1)\right),H_{\alpha}=\int_0^1\frac{1-t^{\alpha}}{1-t}dt=\psi(\alpha+1)+\gamma.$$

Obviously, by a direct calculation, we obtain

$$\zeta_{\alpha}(r)=\zeta(r)-\zeta(r,\alpha+1),\zeta_{n+\alpha}(r)=\zeta_{\alpha}(r)+\sum_{k=1}^{n}\frac{1}{(k+\alpha)^r},$$

where $\zeta(s,a)$ is the Hurwitz zeta function defined by

$$\zeta(r,\alpha+1)=\sum_{n=1}^{\infty}\frac{1}{(n+\alpha)^r},\left(\Re(r)>1,\alpha\notin\mathbb{Z}_0^-,\mathbb{Z}_0^-:=\{0,-1,-2,...\}\right).$$

We now define generalized alternating nth-harmonic number of order r by

$$L_n(r)=\sum_{j=1}^{n}\frac{(-1)^{j-1}}{j^r},\ 1\leq r\in\mathbb{Z}. \tag{1.2}$$

Clearly, we have the following relations for harmonic number and alternating harmonic number

$$L_n(k)=\zeta_{2[(n+1)/2]-1}(k)-\frac{1}{2^k}\left(\zeta_{[n/2]}(k)+\zeta_{[(n-1)/2]}(k)\right),\ 2\leq k\in\mathbb{Z},$$

$$L_n(1)=H_n-H_{[n/2]},$$

where $[x]$ is the integer part of x.
From [19], we know that the classical Euler sums are the infinite sums whose general term is a product of harmonic numbers of index n and a power of n^{-1}. For a pair (p,q) of positive integers with $q\geq 2$, the classical linear Euler sum is defined by

$$S_{p,q}=\sum_{n=1}^{\infty}\frac{\zeta_n(p)}{n^q}. \tag{1.3}$$

The number $w=p+q$ is defined as the weight of $S_{p,q}$. The evaluation of $S_{p,q}$ in terms of values of Riemann zeta function at positive integers is known when $p=1,\ p=q,\ (p,q)=(2,4),(4,2)$

or $p+q$ is odd. In [19], Philippe Flajolet and Bruno Salvy gave a general formula for odd weight $p+q$,

$$\begin{aligned}\sum_{n=1}^{\infty}\frac{\zeta_n(p)}{n^q} &= \zeta(m)\left(\frac{1}{2}-\frac{(-1)^p}{2}\binom{m-1}{p}-\frac{(-1)^p}{2}\binom{m-1}{q}\right)\\ &+(-1)^p\sum_{k=1}^{[p/2]}\binom{m-2k-1}{q-1}\zeta(2k)\zeta(m-2k)+\frac{1-(-1)^p}{2}\zeta(p)\zeta(q)\\ &+(-1)^p\sum_{k=1}^{[q/2]}\binom{m-2k-1}{p-1}\zeta(2k)\zeta(m-2k),\end{aligned}$$

where $\zeta(1)$ should be interpreted as 0 wherever it occurs.
The Riemann zeta function and alternating zeta function are defined respectively by

$$\zeta(s)=\sum_{n=1}^{\infty}\frac{1}{n^s},\Re(s)>1,$$

and

$$\bar{\zeta}(s)=\sum_{n=1}^{\infty}\frac{(-1)^{n-1}}{n^s},\ \Re(s)\geq 1.$$

For a multi-index $S=(s_1,s_2,\cdots,s_k)$ $(s_i\in N,\ s_1\geq 2)$, the general multiple zeta value (MZV for short) and the multiple zeta-star value (MZSV for short) are defined, respectively, by convergent series

$$\zeta(S)=\zeta(s_1,s_2,\cdots,s_k)=\sum_{n_1>n_2>\cdots>n_k\geq 1}\frac{1}{n_1^{s_1}n_2^{s_2}\cdots n_k^{s_k}},$$

$$\zeta^{\star}(S)=\zeta^{\star}(s_1,s_2,\cdots,s_k)=\sum_{n_1\geq n_2\geq\cdots\geq n_k\geq 1}\frac{1}{n_1^{s_1}n_2^{s_2}\cdots n_k^{s_k}}.$$

where $s_1+\cdots+s_k$ is called the weight and k is the multiplicity. Note that our $S_{p,q}$ is called double zeta-star value and denoted by $\zeta^{\star}(q,p)$ in [35]. We put a bar on top of s_j $(j=1,\cdots k)$ if there is a sign $(-1)^{n_j}$ appearing in the denominator on the right. For example,

$$\zeta^{\star}(\bar{2},3)=\sum_{m\geq n\geq 1}\frac{(-1)^m}{m^2n^3}=\sum_{m=1}^{\infty}\frac{(-1)^m}{m^2}\sum_{n=1}^{m}\frac{1}{n^3}.$$

Let $\pi=(\pi_1,\ldots,\pi_k)$ be a partition of integer p and $p=\pi_1+\cdots+\pi_k$ with $\pi_1\leq\pi_2\leq\cdots\leq\pi_k$. The classical nonlinear Euler sum of index π,q is defined as follows (see [19])

$$S_{\pi,q}=\sum_{n=1}^{\infty}\frac{\zeta_n(\pi_1)\zeta_n(\pi_2)\cdots\zeta_n(\pi_k)}{n^q},\tag{1.4}$$

where the quantity $\pi_1+\cdots+\pi_k+q$ is called the weight, the quantity k is called the degree. As usual, repeated summands in partitions are indicated by powers, so that for instance

$$S_{1^2 2^3 4,q}=S_{112224,q}=\sum_{n=1}^{\infty}\frac{H_n^2\zeta_n^3(2)\zeta_n(4)}{n^q}.$$

Euler sums may be studied through a profusion of methods: combinatorial, analytic and algebraic. Many values of nonlinear Euler sums $S_{\pi,q}$ can be expressed as a rational linear combination of zeta values. Explicit evaluations of values of some $S_{\pi,q}$ are presented and proofs in [3, 19, 34]. Table 1 contains some results, some of these identities can be proved by experimental methods, based on [3]. Philippe Flajolet and Bruno Salvy informed us about some ongoing work of theirs ([19]) to evaluate Euler sums in an entirely different way, namely using contour integration and residue theorem. In this way they manage to prove, for instance, that the quadratic sums

$$S_{p_1p_2,q}=\sum_{n=1}^{\infty}\frac{\zeta_n(p_1)\,\zeta_n(p_2)}{n^q}$$

can be evaluated in terms of zeta values and linear sums whenever p_1+p_2+q is even, and $p_1>1, p_2>1, q>1$ (see Theorem 4.2 in the reference [19]]). In [34], we showed that all quadratic Euler sums of the form

$$S_{1m,p}=\sum_{n=1}^{\infty}\frac{H_n\zeta_n(m)}{n^p}\quad(m+p\leq 8)$$

can be reduced to polynomials zeta values and linear sums. The general nonlinear Euler sums involving harmonic number and alternating harmonic number are defined by the series (see [34])

$$\sum_{n=1}^{\infty}\frac{\prod_{i=1}^{m_1}\zeta_n^{q_i}(k_i)\prod_{j=1}^{m_2}L_n^{l_j}(h_j)}{n^p},\ \sum_{n=1}^{\infty}\frac{\prod_{i=1}^{m_1}\zeta_n^{q_i}(k_i)\prod_{j=1}^{m_2}L_n^{l_j}(h_j)}{n^p}(-1)^{n-1},\tag{1.5}$$

where $p(p>1), m_1, m_2, q_i, k_i, h_j, l_j$ are positive integers. If $\sum_{i=1}^{m_1}(k_iq_i)+\sum_{j=1}^{m_2}(h_jl_j)+p=C(C$ is a positive integer), then we call the identity Cth-order Euler sums.
In [19], Philippe Flajolet and Bruno Salvy gave explicit reductions to zeta values and logarithm for all linear sums

$$\sum_{n=1}^{\infty}\frac{\zeta_n(p)}{n^q},\ \sum_{n=1}^{\infty}\frac{L_n(p)}{n^q},\ \sum_{n=1}^{\infty}\frac{\zeta_n(p)}{n^q}(-1)^{n-1},\ \sum_{n=1}^{\infty}\frac{L_n(p)}{n^q}(-1)^{n-1}$$

when $p+q$ is an odd weight. By using the above multiple zeta-star value notations, we have

$$\zeta^{\star}(q,p)=S_{p,q}=\sum_{n=1}^{\infty}\frac{\zeta_n(p)}{n^q},\ \zeta^{\star}(\bar{q},p)=-\sum_{n=1}^{\infty}\frac{\zeta_n(p)}{n^q}(-1)^{n-1},$$

$$\zeta^{\star}(q,\bar{p})=-\sum_{n=1}^{\infty}\frac{L_n(p)}{n^q},\ \zeta^{\star}(\bar{q},\bar{p})=\sum_{n=1}^{\infty}\frac{L_n(p)}{n^q}(-1)^{n-1}.$$

The evaluation of linear sums in terms of values of the Riemann zeta function and polylogarithm function at positive integers is known when $(p,q)=(1,3),(2,2)$, or $p+q$ is odd (see [8, 16, 19, 33]). For example

$$\sum_{n=1}^{\infty}\frac{L_n(1)}{n^3}=\frac{7}{4}\zeta(3)\ln 2-\frac{5}{16}\zeta(4),$$

$$\sum_{n=1}^{\infty}\frac{H_n}{n^3}(-1)^{n-1}=-2\mathrm{Li}_4\left(\frac{1}{2}\right)+\frac{11}{4}\zeta(4)+\frac{1}{2}\zeta(2)\ln^2 2-\frac{1}{12}\ln^4 2-\frac{7}{4}\zeta(3)\ln 2,$$
$$\sum_{n=1}^{\infty}\frac{L_n(1)}{n^3}(-1)^{n-1}=\frac{3}{2}\zeta(4)+\frac{1}{2}\zeta(2)\ln^2 2-\frac{1}{12}\ln^4 2-2\mathrm{Li}_4\left(\frac{1}{2}\right),$$
$$\sum_{n=1}^{\infty}\frac{\zeta_n(2)}{n^2}(-1)^{n-1}=-\frac{51}{16}\zeta(4)+4\mathrm{Li}_4\left(\frac{1}{2}\right)+\frac{7}{2}\zeta(3)\ln 2-\zeta(2)\ln^2(2)+\frac{1}{6}\ln^4 2.$$

Furthermore, we consider the nested sum

$$\sum_{n=1}^{\infty}\frac{y^n}{n^m}\left(\sum_{k=1}^{n}\frac{x^k}{k^p}\right),\ x,y\in[-1,1),\ 1\leq m,p\in\mathbb{Z}.$$

By taking the sum over complementary pairs of summation indices, we obtain a simple reflection formula

$$\sum_{n=1}^{\infty}\frac{y^n}{n^m}\left(\sum_{k=1}^{n}\frac{x^k}{k^p}\right)+\sum_{n=1}^{\infty}\frac{x^n}{n^p}\left(\sum_{k=1}^{n}\frac{y^k}{k^m}\right)=\mathrm{Li}_p(x)\,\mathrm{Li}_m(y)+\mathrm{Li}_{p+m}(xy).\tag{1.6}$$

Hence, we can easily obtain

$$\sum_{n=1}^{\infty}\frac{L_n(2)}{n^2}(-1)^{n-1}=\frac{13}{16}\zeta(4),$$
$$\sum_{n=1}^{\infty}\frac{\zeta_n(3)}{n}(-1)^{n-1}=\frac{19}{16}\zeta(4)-\frac{3}{4}\zeta(3)\ln 2,$$
$$\sum_{n=1}^{\infty}\frac{L_n(2)}{n^2}=\frac{85}{16}\zeta(4)-4\mathrm{Li}_4\left(\frac{1}{2}\right)+\zeta(2)\ln^2 2-\frac{1}{6}\ln^4 2-\frac{7}{2}\zeta(3)\ln 2,$$
$$\sum_{n=1}^{\infty}\frac{L_n(3)}{n}(-1)^{n-1}=2\mathrm{Li}_4\left(\frac{1}{2}\right)+\frac{3}{4}\zeta(3)\ln 2+\frac{1}{12}\ln^4 2-\frac{1}{2}\zeta(4)-\frac{1}{2}\zeta(2)\ln^2 2.$$

In many other cases we are not able to obtain a formula for the Euler sum constant explicitly in terms of values of the Riemann zeta, logarithm and polylogarithm functions, but we are able to obtain relations involving two of more Euler sum constants of the same weight. The polylogarithm function defined for $|x|\leq 1$ by

$$\mathrm{Li}_p(x)=\sum_{n=1}^{\infty}\frac{x^n}{n^p},\Re(p)>1.\tag{1.7}$$

In [35], we proved that the quadratic Euler sums

$$\sum_{n=1}^{\infty}\frac{H_n^2}{n^{2m}}(-1)^{n-1},\sum_{n=1}^{\infty}\frac{L_n^2(1)}{n^{2m}}(-1)^{n-1},\sum_{n=1}^{\infty}\frac{H_nL_n(1)}{n^{2m}}(-1)^{n-1},\sum_{n=1}^{\infty}\frac{H_nL_n(1)}{n^{2m}},\sum_{n=1}^{\infty}\frac{L_n^2(1)}{n^{2m}}$$

are reducible to polynomials zeta values and to linear sums, and are able to give explicit values for certain of these sums in terms of the Riemann zeta function and the polylogarithm function,

m is a positive integer. In [3], David H. Bailey, Jonathan M. Borwein and Roland Girgensohn considered sums

$$\sum_{n=1}^{\infty}\frac{L_n^q(1)}{n^p},\sum_{n=1}^{\infty}\frac{L_n^q(1)}{n^p}(-1)^{n-1},\sum_{n=1}^{\infty}\frac{H_n^q}{n^p}(-1)^{n-1}\ \ (p+q\leq 5)$$

and obtained a number of experimental identities using the experimental method, where p and q are positive integers. Table 2 contains some results (see [3, 35]). The relationship between the values of the Riemann zeta function and Euler sums has been studied by many authors. For explicit evaluations of the classical Euler sums or related alternating sums, readers may consult [1,3,4,8,11,12,16-20,23-35]. The evaluation of harmonic number sums has been useful in various areas of theoretical physics, including in support of Feynman diagram calculations and in resolving open questions on Feynman diagram contributions and relations among special functions [14], including the dilogarithm, Clausen function, and generalized hypergeometric function. An array of harmonic number sums and multiple zeta values is required in calculations of high energy physics. These quantities appear for instance in developing the scattering theory of massless quantum electrodynamics [7]. Harmonic number sums also occur in computer science in the efficiency analysis of algorithms [3].

The main purpose of this paper is to evaluate some Euler-type sums with products of harmonic numbers, Stirling numbers and Bell numbers, either linearly or nonlinearly. Where the Bell numbers $Y_k(n)=Y_k(\zeta_n(1),1!\zeta_n(2),2!\zeta_n(3),\cdots,(r-1)!\zeta_n(r),\cdots)$, $Y_k(x_1,x_2,\cdots)$ stands for the complete exponential Bell polynomial is defined by (see [21])

$$\exp\left(\sum_{m\geq 1}x_m\frac{t^m}{m!}\right)=1+\sum_{k\geq 1}Y_k(x_1,x_2,\cdots)\frac{t^k}{k!}. \tag{1.8}$$

By the definition of the complete exponential Bell polynomial, we have (see [34])

$$Y_0(n)=1,Y_1(n)=H_n,Y_2(n)=H_n^2+\zeta_n(2),Y_3(n)=H_n^3+3H_n\zeta_n(2)+2\zeta_n(3),$$

$$Y_4(n)=H_n^4+8H_n\zeta_n(3)+6H_n^2\zeta_n(2)+3\zeta_n^2(2)+6\zeta_n(4),$$

$$Y_5(n)=H_n^5+10H_n^3\zeta_n(2)+20H_n^2\zeta_n(3)+15H_n\zeta_n^2(2)+30H_n\zeta_n(4)+20\zeta_n(2)\zeta_n(3)+24\zeta_n(5),$$

where $Y_k(n)$ is a rational linear combination of products of harmonic numbers. The plan of the paper ia as follows. In the second section we evaluate some Euler related sums which involve harmonic numbers (or alternating harmonic numbers), Stirling numbers and Bell numbers, and give new closed form sums of several nonlinear Euler sums through Riemann zeta values and polylogarithm functions. As an example we show that

$$\sum_{n=1}^{\infty}\frac{H_n\zeta_n^2(2)}{n^2}=-\frac{217}{16}\zeta(7)+5\zeta(3)\zeta(4)+\frac{13}{2}\zeta(2)\zeta(5).$$

In the third section we use certain integral representations to evaluate several series with harmonic numbers and alternating harmonic numbers. We establish some relationships involving two or more Euler sums of the same weight. In the fourth section, we discuss the analytic representations of triple Euler sums and establish some relations between triple Euler sums and double Euler sums. For instance,

$$\sum_{n=1}^{\infty}\frac{\zeta_n(2)}{n^2}\left(\sum_{k=1}^{n}\frac{H_k^2}{k}\right)=\frac{67}{16}\zeta(7)+\frac{15}{2}\zeta(3)\zeta(4)-\frac{7}{2}\zeta(2)\zeta(5).$$

In the last section we consider the following type of Euler-type sums involving harmonic numbers

$$S_{m_1m_2\ldots m_r}(r,k)=\sum_{n=1}^{\infty}\frac{\zeta_n(m_1)\cdots\zeta_n(m_r)}{(n+r)(n+k)},\ 0\leq r<k\in\mathbb{Z}.$$

We also provide an explicit evaluation of $S_{m_1m_2\ldots m_r}(r,k)$ in a closed form in terms of zeta values and harmonic numbers, and give some several examples. For example

$$\sum_{n=1}^{\infty}\frac{H_n^2}{n(n+k)}=\frac{1}{k}\left\{\begin{array}{l}3\zeta(3)+\dfrac{H_k^3+3H_k\zeta_k(2)+2\zeta_k(3)}{3}\\ -\dfrac{H_k^2+\zeta_k(2)}{k}-\displaystyle\sum_{i=1}^{k-1}\frac{H_i}{i^2}+\zeta(2)H_{k-1}\end{array}\right\}.$$

2 Series involving Stirling numbers, harmonic numbers and Bell numbers

In this section, we will establish some explicit relationships that involve Euler sums and zeta values. The following lemma will be useful in the development of the main theorems.

Lemma 2.1 *(see [34]) Let m,k be integers with $m\geq 2,k\geq 2$. we have*

$$W(m,k)=-\psi^{(m+k)}(1)/(k+1)-\sum_{i=1}^{m-1}\sum_{j=0}^{k-1}\binom{m-1}{i}\binom{k}{j}W(i,j)\psi^{(m+k-i-j-1)}(1),\quad(2.1)$$

where $W(m,k)$ is defined by

$$W(m,k)=\int_0^1\frac{\ln^k(1-x)(\ln x)^m}{1-x}dx.$$

By a direct calculation, we have $W(m,0)=(-1)^m m!\zeta(m+1), W(1,k)=(-1)^{k+1}k!\zeta(k+2)$.

From the recurrence relation (2.1), we know that the values of integrals $W(m,k)$ can be expressed as a rational linear combination of products of zeta values. We now give some values of the integrals $W(m,k)$ for $6\leq m+k\leq 8$

$$\begin{aligned}
&W(5,1)=360\zeta(7)-120\zeta(3)\zeta(4)-120\zeta(2)\zeta(5),\\
&W(4,2)=240\zeta(7)-60\zeta(3)\zeta(4)-96\zeta(2)\zeta(5),\\
&W(3,3)=180\zeta(7)-45\zeta(3)\zeta(4)-72\zeta(2)\zeta(5),\\
&W(4,3)=-\frac{1497}{4}\zeta(8)+576\zeta(3)\zeta(5)-144\zeta(2)\zeta^2(3),\\
&W(3,4)=-366\zeta(8)+432\zeta(3)\zeta(5)-72\zeta(2)\zeta^2(3),\\
&W(4,4)=8064\zeta(9)+288\zeta^3(3)-288\zeta(2)\zeta(7)-1260\zeta(3)\zeta(6)-2016\zeta(4)\zeta(5),\\
&W(3,5)=6720\zeta(9)+120\zeta^3(3)-2160\zeta(2)\zeta(7)-1260\zeta(3)\zeta(6)-1620\zeta(4)\zeta(5).
\end{aligned}$$

$$\sum_{n=1}^{\infty}\frac{H_n\zeta_n(2)}{n^4} = -\frac{51}{16}\zeta(7)+\frac{3}{4}\zeta(3)\zeta(4)+2\zeta(2)\zeta(5),$$

$$\sum_{n=1}^{\infty}\frac{H_n\zeta_n(3)}{n^3} = \frac{83}{8}\zeta(7)+\frac{1}{4}\zeta(3)\zeta(4)-\frac{11}{2}\zeta(2)\zeta(5)$$

$$\sum_{n=1}^{\infty}\frac{H_n\zeta_n(4)}{n^2} = -\frac{51}{16}\zeta(7)-\frac{3}{2}\zeta(3)\zeta(4)+\frac{9}{2}\zeta(2)\zeta(5),$$

$$\sum_{n=1}^{\infty}\frac{\zeta_n^2(2)}{n^3} = -\frac{155}{8}\zeta(7)+\frac{19}{2}\zeta(3)\zeta(4)+5\zeta(2)\zeta(5),$$

$$\sum_{n=1}^{\infty}\frac{\zeta_n(2)\zeta_n(3)}{n^2} = \frac{131}{16}\zeta(7)-\frac{3}{2}\zeta(3)\zeta(4)-\frac{5}{2}\zeta(2)\zeta(5);$$

$$\sum_{n=1}^{\infty}\frac{H_n\zeta_n(2)}{n^5} = -\frac{343}{48}\zeta(8)+12\zeta(3)\zeta(5)-\frac{5}{2}\zeta(2)\zeta^2(3)-\frac{3}{4}S_{2,6},$$

$$\sum_{n=1}^{\infty}\frac{H_n\zeta_n(3)}{n^4} = -\frac{511}{144}\zeta(8)+7\zeta(3)\zeta(5)+\zeta(2)\zeta^2(3)-\frac{25}{4}S_{2,6},$$

$$\sum_{n=1}^{\infty}\frac{H_n\zeta_n(4)}{n^3} = \frac{443}{48}\zeta(8)-\frac{21}{2}\zeta(3)\zeta(5)-\frac{1}{2}\zeta(2)\zeta^2(3)+\frac{25}{4}S_{2,6},$$

$$\sum_{n=1}^{\infty}\frac{H_n\zeta_n(5)}{n^2} = \frac{1063}{144}\zeta(8)-\frac{13}{2}\zeta(3)\zeta(5)+\zeta(2)\zeta^2(3)+\frac{3}{4}S_{2,6},$$

$$\sum_{n=1}^{\infty}\frac{\zeta_n^2(2)}{n^4} = \frac{457}{18}\zeta(8)+6\zeta(2)\zeta^2(3)-40\zeta(3)\zeta(5)+11S_{2,6},$$

$$\sum_{n=1}^{\infty}\frac{\zeta_n^3(2)}{n^2} = \frac{3855}{96}\zeta(8)+9\zeta(2)\zeta^2(3)-60\zeta(3)\zeta(5)+\frac{31}{2}S_{2,6},$$

$$\sum_{n=1}^{\infty}\frac{H_n\zeta_n(2)}{n^6} = \frac{17}{6}\zeta(3)\zeta(6)+\frac{173}{72}\zeta(9)+\frac{1}{4}\zeta(4)\zeta(5)-3\zeta(2)\zeta(7)-\frac{2}{3}\zeta^3(3),$$

$$\sum_{n=1}^{\infty}\frac{H_n\zeta_n(3)}{n^5} = \frac{679}{24}\zeta(9)-11\zeta(2)\zeta(7)-\frac{1}{2}\zeta(3)\zeta(6)-\frac{29}{4}\zeta(4)\zeta(5)-\frac{1}{6}\zeta^3(3),$$

$$\sum_{n=1}^{\infty}\frac{H_n\zeta_n(4)}{n^4} = -\frac{469}{18}\zeta(9)+17\zeta(2)\zeta(7)-\frac{37}{12}\zeta(3)\zeta(6)+2\zeta(4)\zeta(5)+\frac{1}{3}\zeta^3(3),$$

$$\sum_{n=1}^{\infty}\frac{H_n\zeta_n(5)}{n^3} = \frac{679}{24}\zeta(9)-18\zeta(2)\zeta(7)+\frac{3}{4}\zeta(3)\zeta(6)+2\zeta(4)\zeta(5)-\frac{1}{6}\zeta^3(3),$$

$$\sum_{n=1}^{\infty}\frac{H_n\zeta_n(6)}{n^2} = -\frac{799}{72}\zeta(9)+10\zeta(2)\zeta(7)-\frac{5}{2}\zeta(3)\zeta(6)-\frac{1}{2}\zeta(4)\zeta(5)+\frac{1}{3}\zeta^3(3),$$

$$\sum_{n=1}^{\infty}\frac{\zeta_n^2(3)}{n^3} = \frac{253}{6}\zeta(9)+\frac{1}{3}\zeta^3(3)-21\zeta(2)\zeta(7)-6\zeta(4)\zeta(5),$$

$$\sum_{n=1}^{\infty}\frac{\zeta_n^2(2)}{n^5} = -\frac{1069}{36}\zeta(9)+\frac{4}{3}\zeta^3(3)+7\zeta(2)\zeta(7)-\frac{4}{3}\zeta(3)\zeta(6)+\frac{33}{2}\zeta(4)\zeta(5),$$

$$\sum_{n=1}^{\infty}\frac{H_n^2\zeta_n(2)}{n^5} = -\frac{1481}{72}\zeta(9)-3\zeta^3(3)-5\zeta(2)\zeta(7)+\frac{295}{24}\zeta(3)\zeta(6)+18\zeta(4)\zeta(5).$$

TABLE1. Explicit evaluations of nonlinear Euler sums $S_{\pi,q}$

$$\sum_{n=1}^{\infty}\frac{H_nL_n\left(1\right)}{n^2}=\frac{43}{16}\zeta(4)+\frac{3}{4}\zeta(2)\ln^22-\frac{1}{8}\ln^42-3\mathrm{Li}_4\left(\frac{1}{2}\right),$$

$$\sum_{n=1}^{\infty}\frac{L_n^2\left(1\right)}{n^2}=-\frac{13}{8}\zeta(4)+\frac{5}{2}\zeta(2)\ln^22+\frac{1}{12}\ln^42+2\mathrm{Li}_4\left(\frac{1}{2}\right),$$

$$\sum_{n=1}^{\infty}\frac{H_nL_n\left(1\right)}{n^2}(-1)^{n-1}=\frac{29}{16}\zeta\left(4\right)+\frac{3}{4}\zeta\left(2\right)\ln^22-3\mathrm{Li}_4\left(\frac{1}{2}\right)-\frac{1}{8}\ln^42,$$

$$\sum_{n=1}^{\infty}\frac{H_n^2}{n^2}(-1)^{n-1}=\frac{41}{16}\zeta\left(4\right)+\frac{1}{2}\zeta\left(2\right)\ln^22-\frac{1}{12}\ln^42-\frac{7}{4}\zeta\left(3\right)\ln2-2\mathrm{Li}_4\left(\frac{1}{2}\right),$$

$$\sum_{n=1}^{\infty}\frac{L_n^2\left(1\right)}{n^2}(-1)^{n-1}=-\frac{41}{16}\zeta\left(4\right)+2\zeta\left(2\right)\ln^22+\frac{1}{6}\ln^42+\frac{7}{4}\zeta\left(3\right)\ln2+4\mathrm{Li}_4\left(\frac{1}{2}\right),$$

$$\sum_{n=1}^{\infty}\frac{H_n^2}{\left(2n+1\right)^2}=-\frac{61}{16}\zeta\left(4\right)+\frac{7}{2}\zeta\left(2\right)\ln^22+\frac{1}{3}\ln^42+8\mathrm{Li}_4\left(\frac{1}{2}\right),$$

$$\begin{aligned}\sum_{n=1}^{\infty}\frac{H_n^2}{n^3}(-1)^{n-1}&=4\mathrm{Li}_5\left(\frac{1}{2}\right)+4\ln2\mathrm{Li}_4\left(\frac{1}{2}\right)+\frac{2}{15}\ln^52+\frac{7}{4}\zeta\left(3\right)\ln^22-\frac{19}{32}\zeta\left(5\right)\\&\quad-\frac{2}{3}\zeta\left(2\right)\ln^32-\frac{11}{8}\zeta\left(2\right)\zeta\left(3\right),\end{aligned}$$

$$\begin{aligned}\sum_{n=1}^{\infty}\frac{H_n^3}{n^2}(-1)^{n-1}&=6\mathrm{Li}_5\left(\frac{1}{2}\right)+6\ln2\mathrm{Li}_4\left(\frac{1}{2}\right)+\frac{1}{5}\ln^52+\frac{21}{8}\zeta\left(3\right)\ln^22-\frac{9}{4}\zeta\left(5\right)\\&\quad-\zeta\left(2\right)\ln^32-\frac{27}{16}\zeta\left(2\right)\zeta\left(3\right),\end{aligned}$$

$$\sum_{n=1}^{\infty}\frac{H_n^2}{n^4}(-1)^{n-1}=-\frac{175}{32}\zeta\left(6\right)+\frac{1}{2}\zeta^2\left(3\right)+\sum_{n=1}^{\infty}\frac{\zeta_n\left(2\right)}{n^4}(-1)^{n-1}+5\sum_{n=1}^{\infty}\frac{H_n}{n^5}(-1)^{n-1},$$

$$\begin{aligned}\sum_{n=1}^{\infty}\frac{L_n^2\left(1\right)}{n^4}&=\frac{15}{4}\ln^22\zeta\left(4\right)+\frac{9}{4}\zeta\left(2\right)\zeta\left(3\right)\ln2-\frac{31}{4}\zeta\left(5\right)\ln2+\frac{231}{64}\zeta\left(6\right)-\frac{45}{32}\zeta^2\left(3\right)\\&\quad+\sum_{n=1}^{\infty}\frac{L_n\left(1\right)}{n^5}(-1)^{n-1}-\sum_{n=1}^{\infty}\frac{L_n\left(2\right)}{n^4},\end{aligned}$$

$$\begin{aligned}\sum_{n=1}^{\infty}\frac{L_n^2\left(1\right)}{n^4}(-1)^{n-1}&=\frac{15}{4}\ln^22\zeta\left(4\right)+\frac{9}{4}\zeta\left(2\right)\zeta\left(3\right)\ln2-\frac{93}{16}\zeta\left(5\right)\ln2+\frac{35}{64}\zeta\left(6\right)-\frac{15}{16}\zeta^2\left(3\right)\\&\quad+\sum_{n=1}^{\infty}\frac{\zeta_n\left(2\right)}{n^4}(-1)^{n-1},\end{aligned}$$

$$\begin{aligned}\sum_{n=1}^{\infty}\frac{H_nL_n\left(1\right)}{n^4}&=-\frac{3}{2}\zeta\left(2\right)\zeta\left(3\right)\ln2+\frac{31}{8}\zeta\left(5\right)\ln2+\frac{2359}{384}\zeta\left(6\right)-\frac{7}{64}\zeta^2\left(3\right)-\frac{1}{2}\sum_{n=1}^{\infty}\frac{L_n\left(2\right)}{n^4}\\&\quad+\frac{1}{2}\sum_{n=1}^{\infty}\frac{L_n\left(1\right)}{n^5}(-1)^{n-1}-2\sum_{n=1}^{\infty}\frac{\zeta_n\left(2\right)}{n^4}(-1)^{n-1}-4\sum_{n=1}^{\infty}\frac{H_n}{n^5}(-1)^{n-1},\end{aligned}$$

$$\begin{aligned}\sum_{n=1}^{\infty}\frac{H_nL_n\left(1\right)}{n^4}(-1)^{n-1}&=-\frac{3}{2}\zeta\left(2\right)\zeta\left(3\right)\ln2+\frac{155}{32}\zeta\left(5\right)\ln2+\frac{7}{64}\zeta\left(6\right)+\frac{5}{4}\zeta^2\left(3\right)\\&\quad-\sum_{n=1}^{\infty}\frac{\zeta_n\left(2\right)}{n^4}(-1)^{n-1}-\frac{3}{2}\sum_{n=1}^{\infty}\frac{H_n}{n^5}(-1)^{n-1}.\end{aligned}$$

TABLE2. Explicit evaluations of quadratic Euler sums

Lemma 2.2 *(see [34]) For integer $n \geq 1, k \geq 0$, then*

$$\int_0^1 t^{n-1} \ln^k (1-t) dt = (-1)^k \frac{Y_k(n)}{n}, \tag{2.2}$$

where the Bell number $Y_k(n) = Y_k(\zeta_n(1), 1!\zeta_n(2), 2!\zeta_n(3), \cdots, (r-1)!\zeta_n(r), \cdots)$, $Y_k(x_1, x_2, \cdots)$ is defined by (1.8), and $Y_m(n) = m! \sum_{k_1=1}^{n} \frac{1}{k_1} \sum_{k_2=1}^{k_1} \frac{1}{k_2} \cdots \sum_{k_m=1}^{k_{m-1}} \frac{1}{k_m}$.

In fact, applying the definition of Beta function, we know that for $a > 0, 0 < m \in \mathbb{Z}$,

$$\begin{aligned} &\int_0^1 x^{a-1} \ln^m (1-x) dx = \lim_{b \to 1} \frac{\partial^m B(a,b)}{\partial b^m} \\ &= \sum_{i=0}^{m-1} \binom{m-1}{i} \frac{\partial^i B(a,b)}{\partial b^i}\bigg|_{b=1} \left(\psi^{(m-i-1)}(1) - \psi^{(m-i-1)}(a+1) \right), \end{aligned} \tag{2.3}$$

where the Beta function is defined by

$$B(a,b) = \int_0^1 x^{a-1}(1-x)^{b-1} dx = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}, \ \Re(a) > 0, \Re(b) > 0.$$

By using the notation of $\zeta_\alpha(r)$, then (2.3) can be written as

$$\int_0^1 x^{a-1} \ln^m (1-x) dx = \sum_{i=0}^{m-1} (-1)^{m-i} (m-i-1)! \binom{m-1}{i} \left(\int_0^1 x^{a-1} \ln^i (1-x) dx \right) \zeta_a(m-i). \tag{2.4}$$

Therefore, we obtain

$$\begin{aligned} &\int_0^1 x^{a-1} \ln(1-x) dx = -\frac{H_a}{a}, \int_0^1 x^{a-1} \ln^2 (1-x) dx = \frac{H_a^2 + \zeta_a(2)}{a}, \\ &\int_0^1 x^{a-1} \ln^3 (1-x) dx = -\frac{H_a^3 + 3H_a \zeta_a(2) + 2\zeta_a(3)}{a}, \\ &\int_0^1 x^{a-1} \ln^4 (1-x) dx = \frac{H_a^4 + 6H_a^2 \zeta_a(2) + 8H_a \zeta_a(3) + 3\zeta_a^2(2) + 6\zeta_a(4)}{a}. \end{aligned}$$

Define $\frac{Y_m(a)}{a} = (-1)^m \int_0^1 x^{a-1} \ln^m (1-x) dx$, we have the following recurrence relation

$$Y_m(a) = (m-1)! \sum_{i=0}^{m-1} \frac{Y_i(a)}{i!} \zeta_a(m-i), Y_0(a) = 1. \tag{2.5}$$

In general, we will give the following Theorem.

Theorem 2.3 *Define two sequence $A_m(n)$ and $B_m(n)$ by*

$$A_m(n) = (m-1)!\sum_{i=0}^{m-1}\frac{A_i(n)}{i!}\sum_{k=1}^{n}x_k^{m-i}, A_0(n) = 1, (x_k \in \mathbb{R}, k = 1, 2, \cdots, n),$$

$$B_m(n) = \sum_{k_1=1}^{n}x_{k_1}\sum_{k_2=1}^{k_1}x_{k_2}\cdots\sum_{k_m=1}^{k_{m-1}}x_{k_m}, B_0(n) = 1, (x_k \in \mathbb{R}, k = 1, 2, \cdots, n).$$

Then $A_m(n) = m!B_m(n)$.

Proof. By induction, we can tell that $A_m(n)$ and $B_m(n)$ are polynomials of degree m with n variables $x_1, x_2, \cdots x_n$, moreover, $B_m(n)$ has coefficient 1 at $x_1^{s_1}x_2^{s_2}\cdots x_n^{s_n}$, where $\sum_{k=1}^{n}s_k = m$, and $s_k \geq 0$. Suppose $c_{s_1,\cdots,s_n}$ is the coefficient of $A_m(n)$ at $x_1^{s_1}x_2^{s_2}\cdots x_n^{s_n}$, then in order to get the conclusion, all we need to do is prove that $c_{s_1,\cdots,s_n} = m!$ by induction. For convenience, we can suppose $c_{t_1,\cdots,t_n} = 0$ if there is a $t_i < 0$, otherwise $c_{t_1,\cdots,t_n}$ is the coefficient of $A_{t_1+\cdots+t_n}(n)$ at $x_1^{t_1}x_2^{t_2}\cdots x_n^{t_n}$. For $m = 0$, the conclusion holds. If the conclusion holds for any $i \leq m-1$. For m, by the recursion formula of $A_m(n)$, $c_{s_1,\cdots,s_n} = (m-1)!\sum_{i=0}^{m-1}\sum_{k=1}^{n}\frac{c_{s_1,\cdots,s_k-(m-i),\cdots,s_n}}{i!}$, since for $0 \leq i \leq m-1$, $\frac{A_i(n)}{i!}\sum_{k=1}^{n}x_k^{m-i}$ at $x_1^{s_1}x_2^{s_2}\cdots x_n^{s_n} = x_1^{s_1}\cdots x_k^{s_k-(m-i)}\cdots x_n^{s_n}\cdot x_k^{m-i}$, where $1 \leq k \leq n$, provides coefficient $\frac{c_{s_1,\cdots,s_k-(m-i),\cdots,s_n}}{i!}$. So

$$\begin{aligned}
c_{s_1,\cdots,s_n} &= (m-1)!\sum_{k=1}^{n}\sum_{i=0}^{m-1}\frac{c_{s_1,\cdots,s_k-(m-i),\cdots,s_n}}{i!} \\
&= (m-1)!\sum_{k=1}^{n}\sum_{i_k=m-s_k}^{m-1}\frac{c_{s_1,\cdots,s_k-(m-i_k),\cdots,s_n}}{i_k!} \\
&= (m-1)!\sum_{k=1}^{n}\sum_{i_k=m-s_k}^{m-1}1 \\
&= (m-1)!\sum_{k=1}^{n}((m-1)-(m-s_k)+1) \\
&= (m-1)!\sum_{k=1}^{n}s_k \\
&= m!
\end{aligned}$$

since $c_{s_1,\cdots,s_k-(m-i_k),\cdots,s_n} = i_k!$ by the inductive assumption. So the conclusion holds by induction. So far we have completed the proof of Theorem 2.3. □

Let $X_n(m) = \sum_{i=1}^{n}x_i^m$, from Theorem 2.3, we deduce that

$$B_0(n) = 1, B_1(n) = X_n(1), B_2(n) = \frac{X_n^2(1) + X_n(2)}{2},$$

$$B_3(n) = \frac{X_n^3(1) + 3X_n(1)X_n(2) + 2X_n(3)}{3!},$$
$$B_4(n) = \frac{X_n^4(1) + 8X_n(1)X_n(3) + 3X_n^2(2) + 6X_n^2(1)X_n(2) + 6X_n(4)}{4!},$$
$$B_5(n) = \frac{1}{5!}\left\{ \begin{array}{l} X_n^5(1) + 10X_n^3(1)X_n(2) + 20X_n^2(1)X_n(3) + 15X_n(1)X_n^2(2) \\ +30X_n(1)X_n(4) + 20X_n(2)X_n(3) + 24X_n(5) \end{array} \right\},$$
$$B_m(n) = \frac{1}{m}\sum_{i=0}^{m-1} B_i(n)X_n(m-i).$$

Furthermore, we are able to obtain relations involving multiple zeta-star values and zeta values. For example, taking $x_i = \frac{1}{i^p}, p > 1 \ (i = 1, \cdots, n)$ and substituting it into Theorem 2.3, letting $n \to \infty$, we have

$$\zeta^\star(\{p\}^m) = \frac{1}{m}\sum_{i=0}^{m-1} \zeta^\star\left(\{p\}^i\right)\zeta(pm - pi), \tag{2.6}$$

where $\{p\}^m$ denotes the m-tuple $\{p, ..., p\}$, $\zeta^\star\left(\{p\}^0\right) = 1$. Hence ,we obtain

$$\zeta^\star\left(\{2\}^2\right) = \frac{7}{4}\zeta(4), \zeta^\star\left(\{2\}^3\right) = \frac{31}{16}\zeta(6), \zeta^\star\left(\{2\}^4\right) = \frac{127}{64}\zeta(8),$$
$$\zeta^\star\left(\{3\}^2\right) = \frac{1}{2}\left(\zeta^2(3) + \zeta(6)\right), \zeta^\star\left(\{3\}^3\right) = \frac{1}{3!}\left(\zeta^3(3) + 3\zeta(3)\zeta(6) + 2\zeta(9)\right),$$
$$\zeta^\star\left(\{3\}^4\right) = \frac{1}{4!}\left(\zeta^4(3) + 8\zeta(3)\zeta(9) + 3\zeta^2(6) + 6\zeta^2(3)\zeta(6) + 6\zeta(12)\right).$$

By a similar argument as in the proof of Theorem 2.3, we can gives the following Theorem.

Theorem 2.4 *Define two sequence $\bar{A}_m(n)$ and $\bar{B}_m(n)$ by*

$$\bar{A}_m(n) = (m-1)!(-1)^{m-1}\sum_{i=0}^{m-1}(-1)^i\frac{\bar{A}_i(n)}{i!}\sum_{k=1}^{n}x_k^{m-i}, \bar{A}_0(n) = 1,$$

$$\bar{B}_m(n) = \sum_{k_1=1}^{n}x_{k_1}\sum_{k_2=1}^{k_1-1}x_{k_2}\cdots\sum_{k_m=1}^{k_{m-1}-1}x_{k_m}, \bar{B}_0(n) = 1.$$

Then $\bar{A}_m(n) = m!\bar{B}_m(n)$.

Proof. By induction, we can tell that $\bar{A}_m(n)$ and $\bar{B}_m(n)$ are polynomials of degree m with n variables $x_1, x_2, \cdots x_n$, moreover, $\bar{B}_m(n)$ has coefficient 1 at $x_1^{s_1}x_2^{s_2}\cdots x_n^{s_n}$, when $\sum_{k=1}^{n}s_k = m$, and $0 \le s_k \le 1$, otherwise, the coefficient is 0. Suppose $c_{s_1,\cdots,s_n}$ is the coefficient of $\bar{A}_m(n)$ at $x_1^{s_1}x_2^{s_2}\cdots x_n^{s_n}$, then in order to get the conclusion, all we need to do is prove that $c_{s_1,\cdots,s_n} = m!$ if $0 \le s_k \le 1$, and $c_{s_1,\cdots,s_n} = 0$ if there is a k such that $s_k \ge 2$ by induction. For convenience, we can suppose $c_{t_1,\cdots,t_n} = 0$ if there is a $t_i < 0$, otherwise $c_{t_1,\cdots,t_n}$ is the coefficient of $\bar{A}_{t_1+\cdots+t_n}(n)$ at $x_1^{t_1}x_2^{t_2}\cdots x_n^{t_n}$.
For $m = 0$, the conclusion holds. If the conclusion holds for any $i \le m-1$. For m, by the recursion formula of $\bar{A}_m(n)$, $c_{s_1,\cdots,s_n} = (m-1)!(-1)^{m-1}\sum_{i=1}^{m-1}\sum_{k=1}^{n}(-1)^i\frac{c_{s_1,\cdots,s_k-(m-i),\cdots,s_n}}{i!}$, since

for $1 \leq i \leq m-1$, $\frac{\bar{A}_i(n)}{i!}\sum_{k=1}^{n} x_k^{m-i}$ at $x_1^{s_1}x_2^{s_2}\cdots x_n^{s_n} = x_1^{s_1}\cdots x_k^{s_k-(m-i)}\cdots x_n^{s_n}\cdot x_k^{m-i}$, where $1 \leq k \leq n$, provides coefficient $\frac{c_{s_1,\cdots,s_k-(m-i),\cdots,s_n}}{i!}$. So, If there are k_1, k_2 such that $s_{k_1}, s_{k_2} \geq 2$, then $c_{s_1,\cdots,s_k-(m-i),\cdots,s_n} = 0$ for any $0 \leq i \leq m-1, 0 \leq k \leq n$, so $c_{s_1,\cdots,s_n} = 0$. If there is only one k such that $s_k \geq 2$, say j, then $c_{s_1,\cdots,s_k-(m-i),\cdots,s_n} = 0$ for any $0 \leq i \leq m-1, 0 \leq k \leq n$ and $k \neq j$, so

$$\begin{aligned}
c_{s_1,\cdots,s_n} &= (m-1)!(-1)^{m-1}\sum_{i=1}^{m-1}\sum_{k=1}^{n}(-1)^i\frac{c_{s_1,\cdots,s_k-(m-i),\cdots,s_n}}{i!} \\
&= (m-1)!(-1)^{m-1}\sum_{k=1}^{n}\sum_{i_k=m-s_k}^{m-1}(-1)^{i_k}\frac{c_{s_1,\cdots,s_k-(m-i_k),\cdots,s_n}}{i!} \\
&= (m-1)!(-1)^{m-1}\sum_{i=m-s_j}^{m-1}(-1)^i\frac{c_{s_1,\cdots,s_j-(m-i),\cdots,s_n}}{i!} \\
&= (m-1)!(-1)^{m-1}((-1)^{m-s_j}c_{s_1,\cdots,s_{j-1},0,s_{j+1}\cdots,s_n} + (-1)^{m-s_j+1}c_{s_1,\cdots,s_{j-1},1,s_{j+1},\cdots,s_n}) \\
&= (m-1)!(-1)^{m-1}((-1)^{m-s_j} + (-1)^{m-s_j+1}) \\
&= 0
\end{aligned}$$

If $0 \leq s_k \leq 1$ for each $1 \leq k \leq n$, then

$$\begin{aligned}
c_{s_1,\cdots,s_n} &= (m-1)!(-1)^{m-1}\sum_{i=1}^{m-1}\sum_{k=1}^{n}(-1)^i\frac{c_{s_1,\cdots,s_k-(m-i),\cdots,s_n}}{i!} \\
&= (m-1)!(-1)^{m-1}\sum_{k=1}^{n}\sum_{i_k=m-s_k}^{m-1}(-1)^{i_k}\frac{c_{s_1,\cdots,s_k-(m-i_k),\cdots,s_n}}{i_k!} \\
&= (m-1)!(-1)^{m-1}\sum_{\substack{1\leq k\leq n\\ s_k=1}}\sum_{i_k=m-1}^{m-1}(-1)^{i_k}\frac{c_{s_1,\cdots,1-(m-i_k),\cdots,s_n}}{i_k!} \\
&= (m-1)!(-1)^{m-1}\sum_{\substack{1\leq k\leq n\\ s_k=1}}(-1)^{m-1}\frac{c_{s_1,\cdots,s_{k-1},0,s_{k+1},\cdots,s_n}}{(m-1)!} \\
&= (m-1)!\sum_{\substack{1\leq k\leq n\\ s_k=1}}1 \\
&= m!
\end{aligned}$$

Since $c_{s_1,\cdots,s_{k-1},0,s_{k+1},\cdots,s_n} = (\sum_{k=1}^{n} s_k - 1)! = (m-1)!$ by the inductive assumption. So the conclusion holds by induction. □

Let $X_n(m) = \sum_{i=1}^{n} x_i^m$, from Theorem 2.4, we have

$$\bar{B}_1(n) = X_n(1), \bar{B}_2(n) = \frac{X_n^2(1) - X_n(2)}{2!}, \bar{B}_3(n) = \frac{X_n^3(1) - 3X_n(1)X_n(2) + 2X_n(3)}{3!},$$

$$\bar{B}_4(n) = \frac{X_n^4(1) - 6X_n^2(1)X_n(2) + 8X_n(1)X_n(3) + 3X_n^2(2) - 6X_n(4)}{4!},$$
$$\bar{B}_5(n) = \frac{1}{5!}\left\{ \begin{array}{l} X_n^5(1) - 10X_n^3(1)X_n(2) + 20X_n^2(1)X_n(3) + 15X_n(1)X_n^2(2) \\ -30X_n(1)X_n(4) - 20X_n(2)X_n(3) + 24X_n(5) \end{array} \right\},$$
$$\bar{B}_m(n) = \frac{(-1)^{m-1}}{m}\sum_{i=0}^{m-1}(-1)^i \bar{B}_i(n)X_n(m-i).$$

By the definition of $\bar{B}_n(m)$, we can deduce the following relation

$$\bar{B}_m(n-1) = \sum_{i=0}^{m}(-1)^i \bar{B}_{m-i}(n)x_n^i.$$

For instance

$$\begin{aligned}
&\frac{X_{n-1}^4(1) - 6X_{n-1}^2(1)X_{n-1}(2) + 8X_{n-1}(1)X_{n-1}(3) + 3X_{n-1}^2(2) - 6X_{n-1}(4)}{4!} \\
&= \frac{X_n^4(1) - 6X_n^2(1)X_n(2) + 8X_n(1)X_n(3) + 3X_n^2(2) - 6X_n(4)}{4!} \\
&\quad - \frac{X_n^3(1) - 3X_n(1)X_n(2) + 2X_n(3)}{3!}x_n \\
&\quad + \frac{X_n^2(1) - X_n(2)}{2!}x_n^2 - x_n^3X_n(1) + x_n^4, \\
&\frac{X_{n-1}^3(1) - 3X_{n-1}(1)X_{n-1}(2) + 2X_{n-1}(3)}{3!} \\
&= \frac{X_n^3(1) - 3X_n(1)X_n(2) + 2X_n(3)}{3!} \\
&\quad - \frac{X_n^2(1) - X_n(2)}{2!}x_n + x_n^2X_n(1) - x_n^3.
\end{aligned}$$

If $x_n = \frac{1}{n}, \frac{(-1)^{n-1}}{n}$, then we have

$$\begin{aligned}
&\frac{H_{n-1}^4 - 6H_{n-1}^2\zeta_{n-1}(2) + 8H_{n-1}\zeta_{n-1}(3) + 3\zeta_{n-1}^2(2) - 6\zeta_{n-1}(4)}{4!} \\
&= \frac{H_n^4 - 6H_n^2\zeta_n(2) + 8H_n\zeta_n(3) + 3\zeta_n^2(2) - 6\zeta_n(4)}{4!} \\
&\quad - \frac{H_n^3 - 3H_n\zeta_n(2) + 2\zeta_n(3)}{3!n} + \frac{H_n^2 - \zeta_n(2)}{2!n^2} - \frac{H_n}{n^3} + \frac{1}{n^4}, \\
&\frac{H_{n-1}^3 - 3H_{n-1}\zeta_{n-1}(2) + 2\zeta_{n-1}(3)}{3!} \\
&= \frac{H_n^3 - 3H_n\zeta_n(2) + 2\zeta_n(3)}{3!} - \frac{H_n^2 - \zeta_n(2)}{2!n} + \frac{H_n}{n^2} - \frac{1}{n^3}, \\
&\frac{L_{n-1}^4(1) - 6L_{n-1}^2(1)\zeta_{n-1}(2) + 8L_{n-1}(1)L_{n-1}(3) + 3\zeta_{n-1}^2(2) - 6\zeta_{n-1}(4)}{4!} \\
&= \frac{L_n^4(1) - 6L_n^2(1)\zeta_n(2) + 8L_n(1)L_n(3) + 3\zeta_n^2(2) - 6\zeta_n(4)}{4!} \\
&\quad - \frac{L_n^3(1) - 3L_n(1)\zeta_n(2) + 2L_n(3)}{3!n}(-1)^{n-1} + \frac{L_n^2(1) - \zeta_n(2)}{2!n^2} - \frac{L_n(1)}{n^3}(-1)^{n-1} + \frac{1}{n^4},
\end{aligned}$$

$$\frac{L_{n-1}^3(1)-3L_{n-1}(1)\zeta_{n-1}(2)+2L_{n-1}(3)}{3!}$$
$$=\frac{L_n^3(1)-3L_n(1)\zeta_n(2)+2L_n(3)}{3!}-\frac{L_n^2(1)-\zeta_n(2)}{2!n}(-1)^{n-1}+\frac{L_n(1)}{n^2}-\frac{(-1)^{n-1}}{n^3}.$$

Putting $x_i=\frac{1}{i^p}, p>1\ (i=1,\cdots,n)$ and $n\to\infty$ in Theorem 2.4, we obtain

$$\zeta\left(\{p\}^m\right)=\frac{(-1)^{m-1}}{m}\sum_{i=0}^{m-1}(-1)^i\zeta\left(\{p\}^i\right)\zeta(pm-pi). \tag{2.6}'$$

Now we use the identities (2.1) and (2.2) to obtain some identities of Stirling numbers, harmonic numbers and Bell numbers.

Theorem 2.5 *For integers $m,p>0$, we have*

$$\sum_{n=m}^{\infty}\frac{S(n,m)Y_p(n)}{n!n}=p!\binom{m+p}{p}\zeta(m+p+1), \tag{2.7}$$
$$\sum_{n=m}^{\infty}\frac{S(n,m)}{n!n^p}=\sum_{n=p}^{\infty}\frac{S(n,p)}{n!n^m}=\frac{(-1)^{m+p-1}}{m!(p-1)!}W(m,p-1), \tag{2.8}$$
$$\sum_{n=m}^{\infty}\frac{S(n,m)}{n!n}\zeta_n(p+1)=\sum_{n=p}^{\infty}\frac{S(n,p)}{n!n}\zeta_n(m+1), \tag{2.9}$$
$$p!\sum_{n=p}^{\infty}\frac{S(n,p)Y_m(n)}{n!n^2}+m!\sum_{n=m}^{\infty}\frac{S(n,m)Y_p(n)}{n!n^2}=m!p!\zeta(m+1)\zeta(p+1), \tag{2.10}$$

where $S(n,k)$ is called Stirling number of the first kind (see [21, 34]), if $n<k$, then $S(n,k)=0$.

$$S(n,1)=(n-1)!, S(n,2)=(n-1)!H_{n-1},$$
$$S(n,3)=\frac{(n-1)!}{2}\left[H_{n-1}^2-\zeta_{n-1}(2)\right],$$
$$S(n,4)=\frac{(n-1)!}{6}\left[H_{n-1}^3-3H_{n-1}\zeta_{n-1}(2)+2\zeta_{n-1}(3)\right],$$
$$S(n,5)=\frac{(n-1)!}{24}\left[H_{n-1}^4-6\zeta_{n-1}(4)-6H_{n-1}^2\zeta_{n-1}(2)+3\zeta_{n-1}^2(2)+8H_{n-1}\zeta_{n-1}(3)\right].$$

Proof. From [21, 34], we know that

$$\ln^p(1-x)=(-1)^p p!\sum_{n=p}^{\infty}\frac{S(n,p)}{n!}x^n,\ -1\le x<1. \tag{2.11}$$

We therefore have, on using (2.2) and (2.11),

$$\int_0^1\frac{\ln^{p+m}(1-x)}{x}dx=(-1)^m m!\sum_{n=m}^{\infty}\frac{S(n,m)}{n!}\int_0^1 x^{n-1}\ln^p(1-x)\,dx$$
$$=(-1)^{m+p}m!\sum_{n=m}^{\infty}\frac{S(n,m)Y_p(n)}{n!n}. \tag{2.12}$$

The change of variable $x \mapsto 1-x$ now yields

$$\int_0^1 \frac{\ln^{p+m}(1-x)}{x}dx = \int_0^1 \frac{\ln^{p+m}x}{1-x}dx$$
$$= (-1)^{p+m}(p+m)!\zeta(p+m+1). \tag{2.13}$$

Combining (2.12) with (2.13), we can obtain (2.7).
By using (2.11), we have

$$W(m,p-1) = \int_0^1 \frac{\ln^{p-1}(1-x)\ln^m x}{1-x}dx$$
$$= (-1)^{p-1}(p-1)!\sum_{n=p}^{\infty}\frac{S(n,p)}{(n-1)!}\int_0^1 x^{n-1}\ln^m x dx$$
$$= (-1)^{m+p-1}m!(p-1)!\sum_{n=p}^{\infty}\frac{S(n,p)}{n!n^m}$$
$$= \int_0^1 \frac{\ln^{p-1}x\ln^m(1-x)}{x}dx$$
$$= (-1)^m m!\sum_{n=m}^{\infty}\frac{S(n,m)}{n!}\int_0^1 x^{n-1}\ln^{p-1}x dx$$
$$= (-1)^{m+p-1}m!(p-1)!\sum_{n=p}^{\infty}\frac{S(n,m)}{n!n^p}.$$

Hence, we can deduce (2.8). On the other hand, we can easily obtain the following identity

$$\int_0^1 \frac{x^n\ln^m x}{1-x}dx = (-1)^m m!\left(\zeta(m+1)-\zeta_n(m+1)\right),\ 0<m\in\mathbb{Z}. \tag{2.14}$$

Replacing x by t in (2.11), then dividing (2.11) by t and integrating over the interval $(0,x)$, we have

$$\int_0^x \frac{\ln^p(1-t)}{t}dt = (-1)^p p!\sum_{n=p}^{\infty}\frac{S(n,p)}{n!n}x^n, -1\le x\le 1. \tag{2.15}$$

Multiplying (2.15) by $\dfrac{\ln^m x}{1-x}$ and integrating over the interval $(0,1)$, by (2.14), we obtain

$$\int_0^1 \frac{\ln^m x}{1-x}\int_0^x \frac{\ln^p(1-t)}{t}dtdx = (-1)^p p!\sum_{n=p}^{\infty}\frac{S(n,p)}{n!n}\int_0^1 \frac{x^n\ln^m x}{1-x}dx$$
$$= (-1)^{p+m}p!m!\sum_{n=p}^{\infty}\frac{S(n,p)}{n!n}\left(\zeta(m+1)-\zeta_n(m+1)\right). \tag{2.16}$$

Noting that the left-hand side of this equation can be written as

$$\int_0^1 \frac{\ln^m x}{1-x}\int_0^x \frac{\ln^p(1-t)}{t}dtdx \overset{\substack{x=1-u\\ t=1-y}}{=} \int_0^1 \frac{\ln^m(1-u)}{u}\int_u^1 \frac{\ln^p y}{1-y}dydu = \int_0^1 \frac{\ln^m(1-x)}{x}\int_x^1 \frac{\ln^p t}{1-t}dtdx.$$

By integration by parts, we see that

$$\begin{aligned}
&\int_0^1 \frac{\ln^m(1-x)}{x}\int_x^1 \frac{\ln^p t}{1-t}dtdx\\
&= \int_0^1 \left(\int_x^1 \frac{\ln^p t}{1-t}dt\right) d\left(\int_0^x \frac{\ln^m(1-t)}{t}dt\right)\\
&= \left(\int_x^1 \frac{\ln^p t}{1-t}dt\right)\left(\int_0^x \frac{\ln^m(1-t)}{t}dt\right)\Bigg|_0^1 + \int_0^1 \frac{\ln^p x}{1-x}\int_0^x \frac{\ln^m(1-t)}{t}dtdx\\
&= (-1)^{m+p}m!p!\sum_{n=p}^{\infty}\frac{S(n,m)}{n!n}\left(\zeta(p+1)-\zeta_n(p+1)\right). \qquad (2.17)
\end{aligned}$$

Combining (2.15) (2.16) with (2.17), we can obtain (2.9). To prove (2.10), we consider the following integral

$$\begin{aligned}
&\int_0^1\int_0^x \frac{\ln^m(1-x)\ln^p(1-t)}{xt}dtdx\\
&= (-1)^p p!\sum_{n=p}^{\infty}\frac{S(n,p)}{nn!}\int_0^1 x^{n-1}\ln^m(1-x)dx\\
&= (-1)^{p+m}p!\sum_{n=p}^{\infty}\frac{S(n,p)\,Y_m(n)}{n^2n!}\\
&= \left(\int_0^x \frac{\ln^p(1-t)}{t}dt\right)\left(\int_0^x \frac{\ln^m(1-t)}{t}dt\right)\Bigg|_0^1 - \int_0^1\int_0^x \frac{\ln^m(1-t)\ln^p(1-x)}{tx}dtdx\\
&= (-1)^{m+p}m!p!\zeta(m+1)\zeta(p+1) - (-1)^{m+p}m!\sum_{n=m}^{\infty}\frac{S(n,m)\,Y_p(n)}{n^2n!}. \qquad (2.18)
\end{aligned}$$

From (2.18), we can deduce (2.10), we complete the proof of Theorem 2.5. □

Taking $p=2,m=4$ and $p=3,m=3$ in (2.7), $p=2,m=5$ in (2.8), $p=3,m=2$ and $p=4,m=1$ in (2.9), we obtain

$$\sum_{n=1}^{\infty}\frac{\left(H_{n-1}^3-3H_{n-1}\zeta_{n-1}(2)+2\zeta_{n-1}(3)\right)\left(H_n^2+\zeta_n(2)\right)}{n^2} = 180\zeta(7),$$

$$\sum_{n=1}^{\infty}\frac{2H_n^3\zeta_n(2)+2H_n^2\zeta_n(3)-3H_n\zeta_n^2(2)-2\zeta_n(2)\zeta_n(3)}{n^2} - 2\sum_{n=1}^{\infty}\frac{3H_n^2\zeta_n(2)+2H_n\zeta_n(3)}{n^3}$$

$$= \frac{1221}{16}\zeta(7) - 55\zeta(3)\zeta(4) + \frac{11}{2}\zeta(2)\zeta(5),$$

$$\sum_{n=1}^{\infty} \frac{6\zeta_n(4) + 6H_n^2\zeta_n(2) - 3\zeta_n^2(2) - 8H_n\zeta_n(3) - H_n^4}{n^3} = -\frac{W(5,1)}{5} - \frac{W(4,2)}{2!},$$

$$\sum_{n=1}^{\infty} \frac{H_n\zeta_n(3)}{n^3} + \sum_{n=1}^{\infty} \frac{H_n\zeta_n(4)}{n^2} - \frac{1}{2}\sum_{n=1}^{\infty} \frac{H_n^2\zeta_n(3) - \zeta_n(2)\zeta_n(3)}{n^2} = \zeta(3)\zeta(4) + \zeta(7).$$

$$\sum_{n=1}^{\infty} \frac{\left(H_{n-1}^3 - 3H_{n-1}\zeta_{n-1}(2) + 2\zeta_{n-1}(3)\right)\zeta_n(2)}{n^2} = 6\sum_{n=1}^{\infty} \frac{\zeta_n(5)}{n^2}.$$

Noting that, in [3], David H. Bailey, Jonathan M. Borwein and Roland Girgensohn considered sums

$$S_{1^q,p} = \sum_{n=1}^{\infty} \frac{H_n^q}{n^p} \ (p+q = 3,4,5,6,7,8,9)$$

and obtained a number of experimental identities using the experimental method, where p and q are positive integers. From [3, 19], we know that all Euler sums of the form $S_{1^q,p}$ for weights $p+q \in \{3,4,5,6,7,9\}$ are expressible polynomially in terms of zeta values. For weight 8, all such sums are the sum of a polynomial in zeta values and a rational multiple of $S_{2,6}$. For example

$$\sum_{n=1}^{\infty} \frac{H_n^3}{n^4} = \frac{231}{16}\zeta(7) - \frac{51}{4}\zeta(3)\zeta(4) + 2\zeta(2)\zeta(5), \tag{2.19}$$

$$\sum_{n=1}^{\infty} \frac{H_n^4}{n^3} = \frac{185}{8}\zeta(7) - \frac{43}{2}\zeta(3)\zeta(4) + 5\zeta(2)\zeta(5), \tag{2.20}$$

$$\sum_{n=1}^{\infty} \frac{H_n^5}{n^2} = \frac{2051}{16}\zeta(7) + 33\zeta(3)\zeta(4) + \frac{57}{2}\zeta(2)\zeta(5). \tag{2.21}$$

Furthermore, according to (2.19)(2.20) and Table 1, we obtain

$$\sum_{n=1}^{\infty} \frac{H_n^3\zeta_n(2)}{n^2} = \frac{83}{16}\zeta(7) + \frac{27}{2}\zeta(3)\zeta(4) - \frac{5}{2}\zeta(2)\zeta(5),$$

$$\sum_{n=1}^{\infty} \frac{H_n^2\zeta_n(2)}{n^3} = -7\zeta(7) + \frac{19}{2}\zeta(3)\zeta(4) - 2\zeta(2)\zeta(5),$$

$$\sum_{n=1}^{\infty} \frac{H_n^2\zeta_n(3)}{n^2} = \frac{329}{16}\zeta(7) - 6\zeta(3)\zeta(4) - \frac{9}{2}\zeta(2)\zeta(5),$$

$$\sum_{n=1}^{\infty} \frac{H_n\zeta_n^2(2)}{n^2} = -\frac{217}{16}\zeta(7) + 5\zeta(3)\zeta(4) + \frac{13}{2}\zeta(2)\zeta(5).$$

In the same manner, the following identities ar easily derived

$$\sum_{n=1}^{\infty} \frac{H_n\zeta_n(3)}{n^2} = \frac{227}{48}\zeta(6) - \frac{3}{2}\zeta^2(3), \sum_{n=1}^{\infty} \frac{H_n^2\zeta_n(2)}{n^2} = \frac{41}{12}\zeta(6) + 2\zeta^2(3).$$

Therefore, we obtain the following conclusion: If $\sum_{i=1}^{m} k_i q_i + p \leq 7$, then all nonlinear Euler sums

$$\sum_{n=1}^{\infty} \frac{\zeta_n^{q_1}(k_1)\,\zeta_n^{q_2}(k_2)\cdots\zeta_n^{q_m}(k_m)}{n^p}, 2 \leq p \in \mathbb{Z}$$

can be expressed as a rational linear combination of products of Riemann zeta function. Using Mathematica, we can give the numerical values of these nonlinear Euler sums, to 30 decimal digits, see Table 3. Next, we evaluate the following type of Euler sums involving alternating

Euler sums	Closed form	Approx. value
$\sum_{n=1}^{\infty} \frac{H_n^3 \zeta_n(2)}{n^2}$	$\frac{83}{16}\zeta(7) + \frac{27}{2}\zeta(3)\zeta(4) - \frac{5}{2}\zeta(2)\zeta(5)$	18.530307949367856814098562 5627
$\sum_{n=1}^{\infty} \frac{H_n^2 \zeta_n(2)}{n^3}$	$-7\zeta(7) + \frac{19}{2}\zeta(3)\zeta(4) - 2\zeta(2)\zeta(5)$	1.889833567793610785614372 83607
$\sum_{n=1}^{\infty} \frac{H_n^2 \zeta_n(3)}{n^2}$	$\frac{329}{16}\zeta(7) - 6\zeta(3)\zeta(4) - \frac{9}{2}\zeta(2)\zeta(5)$	5.252547277139822199775481 52446
$\sum_{n=1}^{\infty} \frac{H_n \zeta_n^2(2)}{n^2}$	$-\frac{217}{16}\zeta(7) + 5\zeta(3)\zeta(4) + \frac{13}{2}\zeta(2)\zeta(5)$	3.916239128592719894922640 61320

TABLE3. Some numerical values of nonlinear Euler sums

harmonic number, Bell number and Stirling number

$$\sum_{n=m}^{\infty} \frac{L_n(1)\,S(n,m)}{nn!}, \sum_{n=1}^{\infty} \frac{Y_{m-1}(n) L_n(1)}{n}(-1)^{n-1}, \sum_{n=m}^{\infty} \frac{L_n(1)\,S(n,m)}{n!}(-1)^{n-1}.$$

We need the following lemma.

Lemma 2.6 *If $m \geq 1$ is a integer and $z \in [0,1]$, then have*

$$\begin{aligned}\int_0^z \frac{\ln^m(1+x)}{x}dx =& \frac{1}{m+1}\ln^{m+1}(1+z) + m!\left(\zeta(m+1) - \mathrm{Li}_{m+1}\left(\frac{1}{1+z}\right)\right) \\ & - m!\sum_{j=1}^{m} \frac{\ln^{m-j+1}(1+z)}{(m-j+1)!}\mathrm{Li}_j\left(\frac{1}{1+z}\right). \end{aligned} \tag{2.22}$$

Proof. We note that the integral in (2.22), which can be rewritten as

$$\begin{aligned}\int_0^z \frac{\ln^m(1+x)}{x}dx &\overset{t=1+x}{=} \int_1^{1+z} \frac{\ln^m t}{t-1}dt \overset{u=t^{-1}}{=} (-1)^{m+1}\int_1^{(1+z)^{-1}} \frac{\ln^m u}{u-u^2}du \\ &= (-1)^{m+1}\left\{\int_1^{(1+z)^{-1}} \frac{\ln^m u}{u}du + \int_1^{(1+z)^{-1}} \frac{\ln^m u}{1-u}du\right\}\end{aligned}$$

$$=\frac{1}{m+1}\ln^{m+1}(1+z)+(-1)^{m+1}\int_1^{(1+z)^{-1}}\frac{\ln^m u}{1-u}du. \tag{2.23}$$

We use the elementary integral identity

$$\int t^{n-1}(\ln t)^m dt = t^n\left\{\frac{\ln^m t}{n}-\sum_{l=1}^{m}\frac{(-1)^{l+1}\ln^{m-l}(t)(m)_l}{n^{l+1}}\right\}, \tag{2.24}$$

Here $(m)_l = m(m-1)\cdots(m-l+1)$. Then yields

$$\begin{aligned}\int_1^{(1+z)^{-1}}\frac{\ln^m u}{1-u}du=\sum_{n=1}^{\infty}\int_1^{(1+z)^{-1}}u^{n-1}\ln^m u du =&(-1)^{m+1}m!\left(\zeta(m+1)-\mathrm{Li}_{m+1}\left((1+z)^{-1}\right)\right)\\ &+m!(-1)^m\sum_{j=1}^{m}\frac{\ln^{m-j+1}(1+z)}{(m-j+1)!}\mathrm{Li}_j\left((1+z)^{-1}\right).\end{aligned} \tag{2.25}$$

Substituting (2.25) into (2.23) yields the desired result. □

Let $p=3,4$ in (2.22), we get

$$\int_0^1\frac{\ln^3(1+x)}{x}dx=6\zeta(4)+\frac{3}{2}\zeta(2)\ln^2 2-\frac{1}{4}\ln^4 2-\frac{21}{4}\zeta(3)\ln 2-6\mathrm{Li}_4\left(\frac{1}{2}\right),$$

$$\int_0^1\frac{\ln^4(1+x)}{x}dx=-24\mathrm{Li}_5\left(\frac{1}{2}\right)-24\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)-\frac{4}{5}\ln^5 2-\frac{21}{2}\zeta(3)\ln^2 2+24\zeta(5)+4\zeta(2)\ln^3 2.$$

Theorem 2.7 *For integer $m>1$, then have*

$$\int_0^1\frac{\ln^{m+1}(1+x)}{x}dx=\ln 2\int_0^1\frac{\ln^m(1+x)}{x}dx-\ln 2\int_0^1\frac{\ln^m(1-x)}{x}dx+(-1)^m m!\sum_{n=m}^{\infty}\frac{L_n(1)S(n,m)}{nn!}. \tag{2.26}$$

Proof. Using integration by parts we have

$$\int_0^x t^{n-1}\ln(1-t)dt=\frac{1}{n}\left\{x^n\ln(1-x)-\sum_{j=1}^{n}\frac{x^j}{j}-\ln(1-x)\right\}, -1\le x<1. \tag{2.27}$$

Taking $x=-1$ in (2.27), we obtain

$$\int_0^1 x^{n-1}\ln(1+x)dx=\frac{(-1)^n}{n}\left\{(-1)^n\ln 2-\ln 2+L_n(1)\right\}.$$

We consider the following integral

$$\begin{aligned}\int_0^1 \frac{\ln^{m+1}(1+x)}{x}dx &= (-1)^m m!\sum_{n=m}^{\infty} S(n,m)\frac{(-1)^n}{n!}\int_0^1 x^{n-1}\ln(1+x)dx \\ &= (-1)^m m!\sum_{n=m}^{\infty}\frac{S(n,m)}{nn!}\left\{(-1)^n\ln 2-\ln 2+L_n(1)\right\} \\ &= (-1)^m m!\ln 2\sum_{n=m}^{\infty}\frac{S(n,m)}{nn!}(-1)^n-(-1)^m m!\ln 2\sum_{n=m}^{\infty}\frac{S(n,m)}{nn!} \\ &\quad +(-1)^m m!\sum_{n=m}^{\infty}\frac{L_n(1)S(n,m)}{nn!}\end{aligned}$$

Noting that

$$\int_0^1 \frac{\ln^m(1+x)}{x}dx = (-1)^m m!\sum_{n=m}^{\infty}\frac{S(n,m)}{nn!}(-1)^n$$

,

$$\int_0^1 \frac{\ln^m(1-x)}{x}dx = (-1)^m m!\sum_{n=m}^{\infty}\frac{S(n,m)}{nn!},$$

we complete the proof of Theorem 2.7. □

Putting $m=2,3$ in (2.26), we obtain

$$\begin{aligned}&\sum_{n=1}^{\infty}\frac{H_nL_n(1)}{n^2} = \frac{43}{16}\zeta(4)+\frac{3}{4}\zeta(2)\ln^2 2-\frac{1}{8}\ln^4 2-3\mathrm{Li}_4\left(\frac{1}{2}\right), \\ &3\sum_{n=1}^{\infty}\frac{\left(H_n^2-\zeta_n(2)\right)L_n(1)}{n^2}-6\sum_{n=1}^{\infty}\frac{H_nL_n(1)}{n^3}+6\sum_{n=1}^{\infty}\frac{L_n(1)}{n^4} \\ &=12\zeta(4)\ln 2-\frac{5}{2}\zeta(2)\ln^3 2+\frac{11}{20}\ln^5 2+\frac{21}{4}\zeta(3)\ln^2 2+18\mathrm{Li}_4\left(\frac{1}{2}\right)\ln 2+24\mathrm{Li}_5\left(\frac{1}{2}\right)-24\zeta(5).\end{aligned} \tag{2.28}$$

Theorem 2.8 *For integer $m>1$, then have*

$$\sum_{n=1}^{\infty}\frac{Y_{m-1}(n)L_n(1)}{n}(-1)^{n-1} = m!\left(\zeta(m+1)-\mathrm{Li}_{m+1}\left(\frac{1}{2}\right)\right), \tag{2.29}$$

$$\sum_{n=m}^{\infty}\frac{L_n(1)S(n,m)}{n!}(-1)^{n-1} = \mathrm{Li}_{m+1}\left(\frac{1}{2}\right)-\frac{(-1)^m}{m!}\ln^{m+1}2. \tag{2.30}$$

Proof. Firstly, by a direct calculation, we get

$$\int_0^1 \frac{\ln^m(1-x)}{1+x}dx = (-1)^m m!\mathrm{Li}_{m+1}\left(\frac{1}{2}\right). \tag{2.31}$$

On the other hand, by Cauchy product formula, we deduce that

$$\frac{\ln(1-x)}{1+x}=\sum_{n=1}^{\infty}L_n(1)(-x)^n,\ -1<x<1. \tag{2.32}$$

Using (2.2) (2.31) and (2.32), we obtain

$$\begin{aligned}\int_0^1\frac{\ln^m(1-x)}{x(1+x)}dx&=\int_0^1\frac{\ln^m(1-x)}{x}dx-\int_0^1\frac{\ln^m(1-x)}{1+x}dx\\&=(-1)^m m!\left(\zeta(m+1)-\mathrm{Li}_{m+1}\left(\frac{1}{2}\right)\right)\\&=\sum_{n=1}^{\infty}L_n(1)(-1)^n\int_0^1 x^{n-1}\ln^{m-1}(1-x)dx\\&=(-1)^m\sum_{n=1}^{\infty}\frac{Y_{m-1}(n)L_n(1)}{n}(-1)^{n-1}.\end{aligned}$$

Hence, we can obtain (2.29). Secondly, by using (2.11), we have

$$\begin{aligned}\int_0^1\frac{\ln^m(1-x)}{1+x}dx&=(-1)^m m!\sum_{n=m}^{\infty}\frac{S(n,m)}{n!}\int_0^1\frac{x^n}{1+x}dx\\&=(-1)^m m!\sum_{n=m}^{\infty}\frac{S(n,m)}{n!}(-1)^n(\ln 2-L_n(1))\\&=(-1)^m m!\sum_{n=m}^{\infty}\frac{L_n(1)S(n,m)}{n!}(-1)^{n-1}+\ln^{m+1}2.\end{aligned} \tag{2.33}$$

Comparing (2.31) with (2.33) completes the proof of (2.30). □

Remark 2.1 *By applying the known result*

$$\frac{1}{1+x}=\sum_{n=1}^{\infty}(-1)^{n-1}x^{n-1},x\in(-1,1),$$

in (2.31), we obtain

$$\begin{aligned}\int_0^1\frac{\ln^m(1-x)}{1+x}dx&=\sum_{n=1}^{\infty}(-1)^{n-1}\int_0^1 x^{n-1}\ln^m(1-x)dx\\&=(-1)^m\sum_{n=1}^{\infty}\frac{Y_m(n)}{n}(-1)^{n-1}\\&=(-1)^m m!\mathrm{Li}_{m+1}\left(\frac{1}{2}\right).\end{aligned}$$

Therefore we immediately obtain the following result

$$\sum_{n=1}^{\infty}\frac{Y_m(n)}{n}(-1)^{n-1}=m!\mathrm{Li}_{m+1}\left(\frac{1}{2}\right),$$
$$\sum_{n=1}^{\infty}\frac{H_n}{n}(-1)^{n-1}=\frac{\zeta(2)-\ln^2 2}{2},$$
$$\sum_{n=1}^{\infty}\frac{H_n^2+\zeta_n(2)}{n}(-1)^{n-1}=\frac{7}{8}\zeta(3)-\frac{1}{2}\zeta(2)\ln 2+\frac{1}{6}\ln^3 2,$$
$$\sum_{n=1}^{\infty}\frac{H_n^3+3H_n\zeta_n(2)+2\zeta_n(3)}{n}(-1)^{n-1}=6\mathrm{Li}_4\left(\frac{1}{2}\right),$$
$$\sum_{n=1}^{\infty}\frac{H_n^4+8H_n\zeta_n(3)+6H_n^2\zeta_n(2)+3\zeta_n^2(2)+6\zeta_n(4)}{n}(-1)^{n-1}=24\mathrm{Li}_5\left(\frac{1}{2}\right). \tag{2.34}$$

Taking $m=3,4$ in (2.29) (2.30), we get

$$\sum_{n=1}^{\infty}\frac{\left(H_n^2+\zeta_n(2)\right)L_n(1)}{n}(-1)^{n-1}=6\left(\zeta(4)-\mathrm{Li}_4\left(\frac{1}{2}\right)\right), \tag{2.35}$$
$$\sum_{n=1}^{\infty}\frac{\left(H_n^3+3H_n\zeta_n(2)+2\zeta_n(3)\right)L_n(1)}{n}(-1)^{n-1}=24\left(\zeta(5)-\mathrm{Li}_5\left(\frac{1}{2}\right)\right), \tag{2.36}$$
$$\sum_{n=1}^{\infty}\frac{\left(H_n^2-\zeta_n(2)\right)L_n(1)}{n}(-1)^{n-1}-2\sum_{n=1}^{\infty}\frac{H_nL_n(1)}{n^2}(-1)^{n-1}+2\sum_{n=1}^{\infty}\frac{L_n(1)}{n^3}(-1)^{n-1}$$
$$=2\mathrm{Li}_4\left(\frac{1}{2}\right)+\frac{1}{3}\ln^4 2, \tag{2.37}$$
$$\sum_{n=1}^{\infty}\frac{\left(H_n^3-3H_n\zeta_n(2)+\zeta_n(3)\right)L_n(1)}{n}(-1)^{n-1}-3\sum_{n=1}^{\infty}\frac{\left(H_n^2-\zeta_n(2)\right)L_n(1)}{n^2}(-1)^{n-1}$$
$$+6\sum_{n=1}^{\infty}\frac{H_nL_n(1)}{n^3}(-1)^{n-1}-6\sum_{n=1}^{\infty}\frac{L_n(1)}{n^4}(-1)^{n-1}=6\mathrm{Li}_4\left(\frac{1}{2}\right)-\frac{1}{4}\ln^4 2. \tag{2.38}$$

Since $\sum_{n=1}^{\infty}\frac{H_nL_n(1)}{n^2}(-1)^{n-1}$ can be evaluated in terms of zeta values and polylogarithm function (see Table 2), combining (2.35) with (2.37), we give

$$\sum_{n=1}^{\infty}\frac{L_n(1)\zeta_n(2)}{n}(-1)^{n-1}=\frac{43}{16}\zeta(4)-\frac{1}{4}\zeta(2)\ln^2 2-\frac{1}{8}\ln^4 2-3\mathrm{Li}_4\left(\frac{1}{2}\right), \tag{2.39}$$
$$\sum_{n=1}^{\infty}\frac{H_n^2L_n(1)}{n}(-1)^{n-1}=\frac{53}{16}\zeta(4)+\frac{1}{8}\ln^4 2+\frac{1}{4}\zeta(2)\ln^2 2-3\mathrm{Li}_4\left(\frac{1}{2}\right). \tag{2.40}$$

Taking $x_k=\frac{(-1)^{k-1}}{k}$ and $m=4$ in Theorem 2.3, then have

$$X_n(1)=L_n(1),X_n(2)=\zeta_n(2),X_n(3)=L_n(3),X_n(4)=\zeta_n(4).$$

Euler sums	Approx. value
$\sum_{n=1}^{\infty}\frac{H_n^2L_n(1)}{n^2}$	3.42047443750418192202906761725
$\sum_{n=1}^{\infty}\frac{H_n^2L_n(1)}{n^2}(-1)^{n-1}$	1.03612399970649927427318442458
$\sum_{n=1}^{\infty}\frac{L_n(1)\zeta_n(2)}{n^2}$	1.57412834091261541166605196417
$\sum_{n=1}^{\infty}\frac{L_n(1)\zeta_n(2)}{n^2}(-1)^{n-1}$	0.95897256847106732991521801 4827
$\sum_{n=1}^{\infty}\frac{H_nL_n(1)}{n^3}$	1.21543572813973561818275852244
$\sum_{n=1}^{\infty}\frac{H_nL_n(1)}{n^3}(-1)^{n-1}$	0.95587338082626953853595716 0214
$\sum_{n=1}^{\infty}\frac{H_n^3L_n(1)}{n}(-1)^{n-1}$	5.85913107758526146663990336681

TABLE4. The numerical values of 5th-order Euler sums

$$\sum_{k=1}^{n}\frac{L_k^3(1)+3L_k(1)\zeta_k(2)+2L_k(3)}{k}(-1)^{k-1}$$
$$=\frac{1}{4}\left\{L_n^4(1)+8L_n(1)L_n(3)+3\zeta_n^2(2)+6L_n^2(1)\zeta_n(2)+6\zeta_n(4)\right\}. \tag{2.41}$$

Letting $n\to\infty$ in (2.41), we obtain

$$\sum_{n=1}^{\infty}\frac{L_n^3(1)+3L_n(1)\zeta_n(2)+2L_n(3)}{n}(-1)^{n-1}=\frac{1}{4}\ln^4 2+\frac{3}{2}\zeta(3)\ln 2+\frac{27}{8}\zeta(4)+\frac{3}{2}\zeta(2)\ln^2 2. \tag{2.42}$$

Substituting (2.39) into (2.42), we get

$$\sum_{n=1}^{\infty}\frac{L_n^3(1)}{n}(-1)^{n-1}=5\mathrm{Li}_4\left(\frac{1}{2}\right)-\frac{59}{16}\zeta(4)+\frac{13}{4}\zeta(2)\ln^2 2+\frac{11}{24}\ln^4 2. \tag{2.43}$$

In a similar way, we have

$$\sum_{n=1}^{\infty}\frac{L_n^2(1)}{n}(-1)^{n-1}=\frac{1}{3}\ln^3 2+\frac{3}{2}\zeta(2)\ln 2-\frac{1}{2}\zeta(3).$$

By using the experimental method, we can give the numerical values for several Euler sums (see Table 4).

3 Some identities On Euler Sums

In the section, by constructing the integrals of polylogarithm function, we establish some connection involving two or more Euler sums of the same weight. We need the following Theorem.

Theorem 3.1 *Let m,p be integers with $m,p>0$, $|x|,|y|,|z|\in(0,1)$, we have*

$$\begin{aligned}
&\sum_{n=1}^{\infty}\frac{\mathrm{Li}_m(x)\,\zeta_n(p;y)-\mathrm{Li}_p(y)\,\zeta_n(m;x)}{n}z^n\\
&=\mathrm{Li}_p(y)\sum_{n=1}^{\infty}\frac{\zeta_n(1;z)}{n^m}x^n-\mathrm{Li}_m(x)\sum_{n=1}^{\infty}\frac{\zeta_n(1;z)}{n^p}y^n\\
&\quad+\mathrm{Li}_m(x)\,\mathrm{Li}_{p+1}(zy)-\mathrm{Li}_p(y)\,\mathrm{Li}_{m+1}(zx),
\end{aligned}\tag{3.1}$$

where the partial sum $\zeta_n(m;x)$ is defined by $\zeta_n(m;x)=\sum_{k=1}^{n}\frac{x^k}{k^m}$.

Proof. Let us consider the N-th partial sum

$$\begin{aligned}
S_N(x,y,z)&=\sum_{n=1}^{N}\frac{\mathrm{Li}_m(x)\,\zeta_n(p,y)-\mathrm{Li}_p(y)\,\zeta_n(m;x)}{n}z^n\\
&=\mathrm{Li}_m(x)\sum_{n=1}^{N}\frac{\zeta_n(p;y)}{n}z^n-\mathrm{Li}_p(y)\sum_{n=1}^{N}\frac{\zeta_n(m;x)}{n}z^n\\
&=\mathrm{Li}_m(x)\sum_{n=1}^{N}\sum_{k=1}^{n}\frac{y^kz^n}{nk^p}-\mathrm{Li}_p(y)\sum_{n=1}^{N}\sum_{k=1}^{n}\frac{x^kz^n}{nk^m}\\
&=\mathrm{Li}_m(x)\sum_{k=1}^{N}\sum_{n=k}^{N}\frac{y^kz^n}{nk^p}-\mathrm{Li}_p(y)\sum_{n=1}^{N}\sum_{n=k}^{N}\frac{x^kz^n}{nk^m}\\
&=\mathrm{Li}_m(x)\sum_{k=1}^{N}\frac{\zeta_N(1;z)-\zeta_{k-1}(1;z)}{k^p}y^k-\mathrm{Li}_p(y)\sum_{k=1}^{N}\frac{\zeta_N(1;z)-\zeta_{k-1}(1;z)}{k^m}x^k\\
&=\zeta_N(1;z)\left(\mathrm{Li}_m(x)\,\zeta_N(p;y)-\mathrm{Li}_p(y)\,\zeta_N(m;x)\right)\\
&\quad+\mathrm{Li}_p(y)\sum_{k=1}^{N}\frac{\zeta_{k-1}(1;z)}{k^m}x^k-\mathrm{Li}_m(x)\sum_{k=1}^{N}\frac{\zeta_{k-1}(1;z)}{k^p}y^k.
\end{aligned}$$

Letting $N\to\infty$ we can obtain (3.1). □

Taking $(x,y,z)\to(1,1,1),(1,-1,1),(-1,-1,1)$, we get

$$\begin{aligned}
\sum_{n=1}^{\infty}\frac{\zeta(m)\,\zeta_n(p)-\zeta(p)\,\zeta_n(m)}{n}&=\zeta(p)\sum_{n=1}^{\infty}\frac{H_n}{n^m}-\zeta(m)\sum_{n=1}^{\infty}\frac{H_n}{n^p}\\
&\quad+\zeta(m)\,\zeta(p+1)-\zeta(p)\,\zeta(m+1),\quad(m,p\geq2),
\end{aligned}\tag{3.2}$$

$$\begin{aligned}
\sum_{n=1}^{\infty}\frac{\zeta(m)\,L_n(p)-\bar{\zeta}(p)\,\zeta_n(m)}{n}&=\bar{\zeta}(p)\sum_{n=1}^{\infty}\frac{H_n}{n^m}-\zeta(m)\sum_{n=1}^{\infty}\frac{H_n}{n^p}(-1)^{n-1}\\
&\quad+\zeta(m)\,\bar{\zeta}(p+1)-\bar{\zeta}(p)\,\zeta(m+1),\quad(m\geq2,p\geq1),
\end{aligned}\tag{3.3}$$

$$\begin{aligned}
\sum_{n=1}^{\infty}\frac{\bar{\zeta}(m)\,L_n(p)-\bar{\zeta}(p)\,L_n(m)}{n}&=\bar{\zeta}(p)\sum_{n=1}^{\infty}\frac{H_n}{n^m}(-1)^{n-1}-\bar{\zeta}(m)\sum_{n=1}^{\infty}\frac{H_n}{n^p}(-1)^{n-1}\\
&\quad+\bar{\zeta}(m)\,\bar{\zeta}(p+1)-\bar{\zeta}(p)\,\bar{\zeta}(m+1),\quad(m,p\geq1).
\end{aligned}\tag{3.4}$$

The main Theorem of this section can be stated as follows.

Theorem 3.2 *Let $p \geq 2, m \geq 0$ be integers and $x \in [-1,1)$. Then the following identity holds*

$$
\begin{aligned}
&(-1)^{p-1}\sum_{n=1}^{\infty}\left\{\frac{\zeta_n(p+2m+1)}{n^p}+\frac{\zeta_n(p)}{n^{p+2m+1}}\right\}\left(\sum_{k=1}^{n}\frac{x^k}{k}\right)\\
&=\sum_{i=1}^{p+2m}(-1)^{i-1}\mathrm{Li}_{p+2m+2-i}(x)\sum_{n=1}^{\infty}\frac{\zeta_n(p)}{n^i}x^n-\sum_{i=1}^{p-1}(-1)^{i-1}\mathrm{Li}_{p+1-i}(x)\sum_{n=1}^{\infty}\frac{\zeta_n(p+2m+1)}{n^i}x^n\\
&\quad+(-1)^p\ln(1-x)\sum_{n=1}^{\infty}\left\{\frac{\zeta_n(p+2m+1)}{n^p}+\frac{\zeta_n(p)}{n^{p+2m+1}}\right\}(1-x^n). \qquad (3.5)\\
&(-1)^{p-1}\sum_{n=1}^{\infty}\left\{\frac{\zeta_n(p+2m)}{n^p}-\frac{\zeta_n(p)}{n^{p+2m}}\right\}\left(\sum_{k=1}^{n}\frac{x^k}{k}\right)\\
&=\sum_{i=1}^{p+2m-1}(-1)^{i-1}\mathrm{Li}_{p+2m+1-i}(x)\sum_{n=1}^{\infty}\frac{\zeta_n(p)}{n^i}x^n-\sum_{i=1}^{p-1}(-1)^{i-1}\mathrm{Li}_{p+1-i}(x)\sum_{n=1}^{\infty}\frac{\zeta_n(p+2m)}{n^i}x^n\\
&\quad+(-1)^p\ln(1-x)\sum_{n=1}^{\infty}\left\{\frac{\zeta_n(p+2m)}{n^p}-\frac{\zeta_n(p)}{n^{p+2m}}\right\}(1-x^n). \qquad (3.6)
\end{aligned}
$$

Proof. By the definition of polylogarithm function and Cauchy product formula, we can verify that

$$\frac{\mathrm{Li}_m(x)}{1-x}=\sum_{n=1}^{\infty}\zeta_n(m)\,x^n,\ x\in(-1,1). \qquad (3.7)$$

Now, we consider the integral

$$\int_0^x\frac{\mathrm{Li}_p(t)\,\mathrm{Li}_{p+2m+1}(t)}{t(1-t)}dt,\ x\in(-1,1).$$

By virtue of (3.7), we obtain

$$\int_0^x\frac{\mathrm{Li}_p(t)\,\mathrm{Li}_{p+2m+1}(t)}{t(1-t)}dt=\sum_{n=1}^{\infty}\zeta_n(p)\int_0^x t^{n-1}\mathrm{Li}_{p+2m+1}(t)dt=\sum_{n=1}^{\infty}\zeta_n(p+2m+1)\int_0^x t^{n-1}\mathrm{Li}_p(t)dt. \qquad (3.8)$$

On the other hand, using integration by parts we have

$$\int_0^x t^{n-1}\mathrm{Li}_q(t)\,dt=\sum_{i=1}^{q-1}(-1)^{i-1}\frac{x^n}{n^i}\mathrm{Li}_{q+1-i}(x)+\frac{(-1)^q}{n^q}\ln(1-x)(x^n-1)-\frac{(-1)^q}{n^q}\left(\sum_{k=1}^{n}\frac{x^k}{k}\right). \qquad (3.9)$$

Substituting (3.9) into (3.8) respectively, we get

$$
\begin{aligned}
&\int_0^x\frac{\mathrm{Li}_p(t)\,\mathrm{Li}_{p+2m+1}(t)}{t(1-t)}dt\\
&=\sum_{i=1}^{p+2m}(-1)^{i-1}\mathrm{Li}_{p+2m+2-i}(x)\sum_{n=1}^{\infty}\frac{\zeta_n(p)}{n^i}x^n+(-1)^p\ln(1-x)\sum_{n=1}^{\infty}\frac{\zeta_n(p)}{n^{p+2m+1}}(1-x^n)
\end{aligned}
$$

$$+(-1)^p\sum_{n=1}^{\infty}\frac{\zeta_n(p)}{n^{p+2m+1}}\left(\sum_{k=1}^{n}\frac{x^k}{k}\right)$$
$$=\sum_{i=1}^{p-1}(-1)^{i-1}\mathrm{Li}_{p+1-i}(x)\sum_{n=1}^{\infty}\frac{\zeta_n(p+2m+1)}{n^i}x^n+(-1)^{p-1}\ln(1-x)\sum_{n=1}^{\infty}\frac{\zeta_n(p+2m+1)}{n^p}(1-x^n)$$
$$+(-1)^{p-1}\sum_{n=1}^{\infty}\frac{\zeta_n(p+2m+1)}{n^p}\left(\sum_{k=1}^{n}\frac{x^k}{k}\right). \tag{3.10}$$

By a direct calculation, we obtain (3.5) holds. Similarly to the proof of (3.5), considering integral

$$\int_0^x\frac{\mathrm{Li}_p(t)\,\mathrm{Li}_{p+2m}(t)}{t(1-t)}dt,\ x\in(-1,1),$$

we deduce (3.6) holds. □

Proceeding in a similar fashion to evaluation of the Theorem 2.2, we can give the following Theorems.

Theorem 3.3 *Let $p\geq 1, m\geq 2$ be integers and $x\in(-1,1)$, then we have*

$$(-1)^m\sum_{n=1}^{\infty}\frac{L_n(p)}{n^m}\left(\sum_{k=1}^{n}\frac{x^k}{k}\right)-(-1)^p\sum_{n=1}^{\infty}\frac{\zeta_n(m)}{n^p}\left(\sum_{k=1}^{n}\frac{(-x)^k}{k}\right)(-1)^{n-1}$$
$$=\sum_{i=1}^{p-1}(-1)^{i-1}\mathrm{Li}_{p+1-i}(-x)\sum_{n=1}^{\infty}\frac{\zeta_n(m)}{n^i}x^n+\sum_{i=1}^{m-1}(-1)^{i-1}\mathrm{Li}_{m+1-i}(x)\sum_{n=1}^{\infty}\frac{L_n(p)}{n^i}x^n$$
$$+(-1)^p\ln(1+x)\sum_{n=1}^{\infty}\frac{\zeta_n(m)}{n^p}\left(x^n+(-1)^{n-1}\right)+(-1)^m\ln(1-x)\sum_{n=1}^{\infty}\frac{L_n(p)}{n^m}(x^n-1), \tag{3.11}$$
$$(-1)^m\sum_{n=1}^{\infty}\frac{L_n(p)}{n^m}\left(\sum_{k=1}^{n}\frac{x^k}{k}\right)(-1)^{n-1}-(-1)^p\sum_{n=1}^{\infty}\frac{L_n(m)}{n^p}\left(\sum_{k=1}^{n}\frac{x^k}{k}\right)(-1)^{n-1}$$
$$=\sum_{i=1}^{m-1}(-1)^{i-1}\mathrm{Li}_{m+1-i}(x)\sum_{n=1}^{\infty}\frac{L_n(p)}{n^i}x^n(-1)^{n-1}-\sum_{i=1}^{p-1}(-1)^{i-1}\mathrm{Li}_{p+1-i}(x)\sum_{n=1}^{\infty}\frac{L_n(m)}{n^i}x^n(-1)^{n-1}$$
$$+(-1)^m\ln(1-x)\sum_{n=1}^{\infty}\frac{L_n(p)}{n^m}(-1)^{n-1}(x^n-1)-(-1)^p\ln(1-x)\sum_{n=1}^{\infty}\frac{L_n(m)}{n^p}(-1)^{n-1}(x^n-1). \tag{3.12}$$

Proof. Proceeding in the similar way, we consider the integrals

$$\int_0^x\frac{\mathrm{Li}_m(t)\,\mathrm{Li}_p(-t)}{(1-t)t}dt,\int_0^x\frac{\mathrm{Li}_m(t)\,\mathrm{Li}_p(t)}{(1+t)t}dt.$$

By a direct computation, we can deduce Theorem 3.3. □

Taking $x\to 1$ in (3.11), $x\to -1$ in (3.12) and using (3.3) (3.4), we can gives the following Corollaries.

Corollary 3.4 *For $p\geq 2, m\geq 2$ and $p,m\in\mathbb{Z}$, we have*

$$\begin{aligned}
&(-1)^m\sum_{n=1}^{\infty}\frac{H_nL_n\left(p\right)}{n^m}+(-1)^p\sum_{n=1}^{\infty}\frac{L_n\left(1\right)\zeta_n\left(m\right)}{n^p}(-1)^{n-1}\\
&=\sum_{i=2}^{m-1}(-1)^{i-1}\zeta\left(m+1-i\right)\sum_{n=1}^{\infty}\frac{L_n\left(p\right)}{n^i}-\sum_{i=2}^{p-1}(-1)^{i-1}\bar{\zeta}\left(p+1-i\right)\sum_{n=1}^{\infty}\frac{\zeta_n\left(m\right)}{n^i}\\
&\quad+\bar{\zeta}\left(p\right)\sum_{n=1}^{\infty}\frac{H_n}{n^m}-\zeta\left(m\right)\sum_{n=1}^{\infty}\frac{H_n}{n^p}(-1)^{n-1}\ +\zeta\left(m\right)\bar{\zeta}\left(p+1\right)-\bar{\zeta}\left(p\right)\zeta\left(m+1\right)\\
&\quad+(-1)^p\ln 2\sum_{n=1}^{\infty}\frac{\zeta_n\left(m\right)}{n^p}\left(1+(-1)^{n-1}\right).
\end{aligned}\tag{3.13}$$

Corollary 3.5 *For $p\geq 2, m\geq 2$ and $p,m\in\mathbb{Z}$, we have*

$$\begin{aligned}
&(-1)^p\sum_{n=1}^{\infty}\frac{L_n\left(1\right)L_n\left(m\right)}{n^p}(-1)^{n-1}-(-1)^m\sum_{n=1}^{\infty}\frac{L_n\left(1\right)L_n\left(p\right)}{n^m}(-1)^{n-1}\\
&=\sum_{i=2}^{m-1}(-1)^{i-1}\bar{\zeta}\left(m+1-i\right)\sum_{n=1}^{\infty}\frac{L_n\left(p\right)}{n^i}-\sum_{i=2}^{p-1}(-1)^{i-1}\bar{\zeta}\left(p+1-i\right)\sum_{n=1}^{\infty}\frac{L_n\left(m\right)}{n^i}\\
&+\bar{\zeta}\left(p\right)\sum_{n=1}^{\infty}\frac{H_n}{n^m}(-1)^{n-1}-\bar{\zeta}\left(m\right)\sum_{n=1}^{\infty}\frac{H_n}{n^p}(-1)^{n-1}+\bar{\zeta}\left(m\right)\bar{\zeta}\left(p+1\right)-\bar{\zeta}\left(p\right)\bar{\zeta}\left(m+1\right)\\
&+(-1)^p\ln 2\sum_{n=1}^{\infty}\frac{L_n\left(m\right)}{n^p}\left(1+(-1)^{n-1}\right)-(-1)^m\ln 2\sum_{n=1}^{\infty}\frac{L_n\left(p\right)}{n^m}\left(1+(-1)^{n-1}\right).
\end{aligned}\tag{3.14}$$

Similarly, taking $x\to -1, p=1$ in (3.11) and $x\to 1, p=1$ in (3.12), we obtain the following Corollaries.

Corollary 3.6 *For integer $m>1$, then*

$$\begin{aligned}
&(-1)^{m-1}\sum_{n=1}^{\infty}\frac{L_n^2\left(1\right)}{n^m}+\sum_{n=1}^{\infty}\frac{H_n\zeta_n\left(m\right)}{n}(-1)^{n-1}\\
&=\sum_{i=1}^{m-1}(-1)^{i-1}\bar{\zeta}\left(m+1-i\right)\sum_{n=1}^{\infty}\frac{L_n\left(1\right)}{n^i}(-1)^{n-1}+(-1)^{m-1}\ln 2\sum_{n=1}^{\infty}\frac{L_n\left(1\right)}{n^m}\left(1+(-1)^{n-1}\right).
\end{aligned}\tag{3.15}$$

Corollary 3.7 *For integer $m>1$, then*

$$\begin{aligned}
&(-1)^m\sum_{n=1}^{\infty}\frac{H_nL_n\left(1\right)}{n^m}(-1)^{n-1}+\sum_{n=1}^{\infty}\frac{H_nL_n\left(m\right)}{n}(-1)^{n-1}\\
&=\sum_{i=1}^{m-1}(-1)^{i-1}\zeta\left(m+1-i\right)\sum_{n=1}^{\infty}\frac{L_n\left(1\right)}{n^i}(-1)^{n-1}.
\end{aligned}\tag{3.16}$$

From (3.11)-(3.16), we give some example:

$$\sum_{n=1}^{\infty}\frac{L_n\left(1\right)L_n\left(2\right)}{n^2}+\sum_{n=1}^{\infty}\frac{H_n\zeta_n\left(2\right)}{n^2}(-1)^{n-1}=-4\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)-\frac{1}{6}\ln^5 2-\frac{7}{2}\zeta\left(3\right)\ln^2 2$$

$$+\zeta(2)\ln^3 2+\frac{3}{16}\zeta(2)\zeta(3)+\frac{49}{8}\zeta(4)\ln 2, \quad (3.17)$$

$$\sum_{n=1}^{\infty}\frac{H_nL_n(2)}{n^2}+\sum_{n=1}^{\infty}\frac{L_n(1)\zeta_n(2)}{n^2}(-1)^{n-1}=4\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)+\frac{1}{6}\ln^5 2+\frac{7}{2}\zeta(3)\ln^2 2$$
$$-\zeta(2)\ln^3 2+\frac{5}{8}\zeta(2)\zeta(3)-\frac{23}{16}\zeta(4)\ln 2, \quad (3.18)$$

$$\sum_{n=1}^{\infty}\frac{H_n\zeta_n(2)}{n}(-1)^{n-1}-\sum_{n=1}^{\infty}\frac{L_n^2(1)}{n^2}=\frac{5}{8}\zeta(4)+\frac{7}{8}\zeta(3)\ln 2-\frac{11}{4}\zeta(2)\ln^2 2, \quad (3.19)$$

$$\sum_{n=1}^{\infty}\frac{H_n\zeta_n(3)}{n}(-1)^{n-1}+\sum_{n=1}^{\infty}\frac{L_n^2(1)}{n^3}=-2\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)-\frac{1}{12}\ln^5 2+\frac{17}{8}\zeta(3)\ln^2 2,$$
$$+\frac{1}{2}\zeta(2)\ln^3 2+\frac{11}{16}\zeta(2)\zeta(3)-\frac{11}{16}\zeta(4)\ln 2, \quad (3.20)$$

$$\sum_{n=1}^{\infty}\frac{H_nL_n(1)}{n^2}(-1)^{n-1}+\sum_{n=1}^{\infty}\frac{H_nL_n(2)}{n}(-1)^{n-1}=\frac{5}{4}\zeta(4)+\frac{1}{2}\zeta(2)\ln^2 2, \quad (3.21)$$

$$\sum_{n=1}^{\infty}\frac{H_nL_n(3)}{n}(-1)^{n-1}-\sum_{n=1}^{\infty}\frac{H_nL_n(1)}{n^3}(-1)^{n-1}=\frac{1}{2}\zeta(3)\ln^2 2+\frac{9}{8}\zeta(2)\zeta(3)-\frac{15}{4}\zeta(4)\ln 2. \quad (3.22)$$

Combining (3.19) (3.21) with Table 2, we obtain

$$\sum_{n=1}^{\infty}\frac{H_nL_n(2)}{n}(-1)^{n-1}=3\mathrm{Li}_4\left(\frac{1}{2}\right)+\frac{1}{8}\ln^4 2-\frac{1}{4}\zeta(2)\ln^2 2-\frac{9}{16}\zeta(4), \quad (3.23)$$

$$\sum_{n=1}^{\infty}\frac{H_n\zeta_n(2)}{n}(-1)^{n-1}=2\mathrm{Li}_4\left(\frac{1}{2}\right)+\frac{1}{12}\ln^4 2+\frac{7}{8}\zeta(3)\ln 2-\frac{1}{4}\zeta(2)\ln^2 2-\zeta(4). \quad (3.24)$$

In (2.11), taking $x=-1, p=4,5$, we have

$$\sum_{n=1}^{\infty}\frac{H_n^3}{n+1}(-1)^{n-1}-3\sum_{n=1}^{\infty}\frac{H_n\zeta_n(2)}{n+1}(-1)^{n-1}+2\sum_{n=1}^{\infty}\frac{\zeta_n(3)}{n+1}(-1)^{n-1}=\frac{1}{4}\ln^4 2, \quad (3.25)$$

$$\sum_{n=1}^{\infty}\frac{H_n^4+8H_n\zeta_n(3)-6H_n^2\zeta_n(2)+3\zeta_n^2(2)-6\zeta_n(4)}{n}(-1)^{n-1}=\frac{1}{5}\ln^5 2-\int_0^1\frac{\ln^4(1+x)}{x}dx. \quad (3.25)'$$

Substituting related equations into (3.25), we get

$$\sum_{n=1}^{\infty}\frac{H_n^3}{n}(-1)^{n-1}=\frac{5}{8}\zeta(4)+\frac{3}{4}\zeta(2)\ln^2 2-\frac{1}{4}\ln^4 2-\frac{9}{8}\zeta(3)\ln 2. \quad (3.26)$$

This formula was also given by I.Mezö in [24]. From (2.22)(2.34) and (3.25)′, we also obtain

$$\sum_{n=1}^{\infty}\frac{H_n^2\zeta_n(2)}{n}(-1)^{n-1}=-2\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)-\frac{1}{12}\ln^5 2-\frac{7}{8}\zeta(3)\ln^2 2+\frac{1}{3}\zeta(2)\ln^3 2$$
$$+\frac{3}{8}\zeta(2)\zeta(3)+\frac{7}{8}\zeta(4)\ln 2.$$

In [35], we gave the following formula: for $s \geq 2$,

$$\frac{1}{2}\sum_{n=1}^{\infty}\frac{H_n^2}{n^s}(-1)^{n-1}-\sum_{n=1}^{\infty}\frac{H_nL_n\left(1\right)}{n^s}(-1)^{n-1}=-\frac{3}{2}\zeta^{\star}\left(\overline{s},2\right)-s\zeta^{\star}\left(\overline{s+1},1\right)-\zeta^{\star}\left(s+1,1\right)$$
$$-\sum_{j=2}^{s}\zeta^{\star}\left(j,1\right)\bar{\zeta}\left(s+1-j\right)+\ln 2\zeta^{\star}\left(\overline{s},1\right), \quad (3.27)$$

From Table 3 in the reference [3], we can obtain

$$\sum_{n=1}^{\infty}\frac{H_n^3}{n^2}(-1)^{n-1}=6\mathrm{Li}_5\left(\frac{1}{2}\right)+6\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)+\frac{1}{5}\ln^5 2+\frac{21}{8}\zeta\left(3\right)\ln^2 2$$
$$-\frac{9}{4}\zeta\left(5\right)-\zeta\left(2\right)\ln^3 2-\frac{27}{16}\zeta\left(2\right)\zeta\left(3\right), \quad (3.28)$$
$$\sum_{n=1}^{\infty}\frac{H_n^2}{n^3}(-1)^{n-1}=4\mathrm{Li}_5\left(\frac{1}{2}\right)+4\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)+\frac{2}{15}\ln^5 2+\frac{7}{4}\zeta\left(3\right)\ln^2 2$$
$$-\frac{19}{32}\zeta\left(5\right)-\frac{2}{3}\zeta\left(2\right)\ln^3 2-\frac{11}{8}\zeta\left(2\right)\zeta\left(3\right), \quad (3.29)$$
$$\sum_{n=1}^{\infty}\frac{L_n^2\left(1\right)}{n^3}=4\mathrm{Li}_5\left(\frac{1}{2}\right)-\frac{1}{30}\ln^5 2+\frac{7}{4}\zeta\left(3\right)\ln^2 2-\frac{167}{32}\zeta\left(5\right)+\frac{1}{3}\zeta\left(2\right)\ln^3 2$$
$$+\frac{3}{4}\zeta\left(2\right)\zeta\left(3\right)+\frac{19}{8}\zeta\left(4\right)\ln 2, \quad (3.30)$$
$$\sum_{n=1}^{\infty}\frac{L_n^2\left(1\right)}{n^3}(-1)^{n-1}=-4\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)-\frac{1}{6}\ln^5 2-\frac{19}{32}\zeta\left(5\right)+\zeta\left(2\right)\ln^3 2$$
$$+\frac{3}{8}\zeta\left(2\right)\zeta\left(3\right)+\frac{19}{8}\zeta\left(4\right)\ln 2. \quad (3.31)$$

Applying (3.30) to (3.20), we deduce that

$$\sum_{n=1}^{\infty}\frac{H_n\zeta_n\left(3\right)}{n}(-1)^{n-1}=-4\mathrm{Li}_5\left(\frac{1}{2}\right)-2\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)-\frac{1}{20}\ln^5 2+\frac{3}{8}\zeta\left(3\right)\ln^2 2+\frac{167}{32}\zeta\left(5\right)$$
$$+\frac{1}{6}\zeta\left(2\right)\ln^3 2-\frac{1}{16}\zeta\left(2\right)\zeta\left(3\right)-\frac{49}{16}\zeta\left(4\right)\ln 2. \quad (3.32)$$

Taking $s=3$ in (3.27), then substituting (3.29) into (3.25), we have

$$\sum_{n=1}^{\infty}\frac{H_nL_n\left(1\right)}{n^3}(-1)^{n-1}=2\mathrm{Li}_5\left(\frac{1}{2}\right)-\frac{1}{60}\ln^5 2-\frac{37}{16}\zeta\left(5\right)-\frac{7}{8}\zeta\left(3\right)\ln^2 2$$
$$+\frac{1}{6}\zeta\left(2\right)\ln^3 2+4\zeta\left(4\right)\ln 2-\frac{1}{8}\zeta\left(2\right)\zeta\left(3\right). \quad (3.33)$$

Substituting (3.33) into (3.22) respectively, the result is

$$\sum_{n=1}^{\infty}\frac{H_nL_n\left(3\right)}{n}(-1)^{n-1}=2\mathrm{Li}_5\left(\frac{1}{2}\right)-\frac{1}{60}\ln^5 2-\frac{3}{8}\zeta\left(3\right)\ln^2 2-\frac{37}{16}\zeta\left(5\right)+\frac{1}{6}\zeta\left(2\right)\ln^3 2$$
$$+\zeta\left(2\right)\zeta\left(3\right)+\frac{1}{4}\zeta\left(4\right)\ln 2. \quad (3.34)$$

Let $m=4$ in Lemma 2.5, we arrive at the conclusion that

$$\begin{aligned}\int_0^1 \frac{\ln^4(1+x)}{x}dx &= 4\sum_{n=1}^{\infty}\frac{H_n^3-3H_n\zeta_n(2)+2\zeta_n(3)}{(n+1)^2}(-1)^{n+1}\\ &= -24\mathrm{Li}_5\left(\frac{1}{2}\right)-24\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)-\frac{4}{5}\ln^5 2-\frac{21}{2}\zeta(3)\ln^2 2\\ &\quad+24\zeta(5)+4\zeta(2)\ln^3 2. \end{aligned} \tag{3.35}$$

After a little simplification, we deduce that

$$\begin{aligned}\sum_{n=1}^{\infty}\frac{H_n\zeta_n(2)}{n^2}(-1)^{n-1} &= -4\mathrm{Li}_5\left(\frac{1}{2}\right)-4\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)-\frac{2}{15}\ln^5 2-\frac{7}{4}\zeta(3)\ln^2 2\\ &\quad+\frac{23}{8}\zeta(5)+\frac{2}{3}\zeta(2)\ln^3 2+\frac{15}{16}\zeta(2)\zeta(3). \end{aligned} \tag{3.36}$$

Combining (3.17) with (3.36) gives

$$\begin{aligned}\sum_{n=1}^{\infty}\frac{L_n(1)L_n(2)}{n^2} &= 4\mathrm{Li}_5\left(\frac{1}{2}\right)-\frac{1}{30}\ln^5 2-\frac{7}{4}\zeta(3)\ln^2 2-\frac{23}{8}\zeta(5)+\frac{1}{3}\zeta(2)\ln^3 2\\ &\quad-\frac{3}{4}\zeta(2)\zeta(3)+\frac{49}{8}\zeta(4)\ln 2. \end{aligned} \tag{3.37}$$

In [35], we proved that the quadratic Euler sums

$$\sum_{n=1}^{\infty}\frac{H_nL_n(1)}{n^{2m}}(-1)^{n-1},\sum_{n=1}^{\infty}\frac{L_n^2(1)}{n^{2m}}$$

are reducible to polynomials zeta values and to linear sums. Hence, from Corollary 3.6 and Corollary 3.7, we obtain the following Theorem.

Theorem 3.8 *For integer $m>1$, the quadratic sums*

$$\sum_{n=1}^{\infty}\frac{H_n\zeta_n(2m)}{n}(-1)^{n-1},\sum_{n=1}^{\infty}\frac{H_nL_n(2m)}{n}(-1)^{n-1}$$

are reducible to linear sums.

Using (3.15) (3.16) and Table 2, we can derive

$$\begin{aligned}\sum_{n=1}^{\infty}\frac{H_n\zeta_n(4)}{n}(-1)^{n-1} &= \frac{105}{32}\zeta(6)-\frac{93}{32}\zeta(5)\ln 2-\frac{3}{16}\zeta^2(3)+\frac{17}{16}\zeta(4)\ln^2 2-\frac{1}{24}\zeta(2)\ln^4 2\\ &\quad-\zeta(2)\mathrm{Li}_4\left(\frac{1}{2}\right)+\sum_{n=1}^{\infty}\frac{H_n}{n^5}(-1)^{n-1}-\sum_{n=1}^{\infty}\frac{L_n(2)}{n^4}, \end{aligned} \tag{3.38}$$

$$\begin{aligned}\sum_{n=1}^{\infty}\frac{H_nL_n(4)}{n}(-1)^{n-1} &= \frac{1}{8}\zeta^2(3)+\frac{155}{32}\zeta(5)\ln 2+\frac{441}{64}\zeta(6)+\frac{7}{4}\zeta(4)\ln^2 2-\frac{1}{12}\zeta(2)\ln^4 2\\ &\quad-2\zeta(2)\mathrm{Li}_4\left(\frac{1}{2}\right)-\sum_{n=1}^{\infty}\frac{\zeta_n(2)}{n^4}(-1)^{n-1}-\frac{3}{2}\sum_{n=1}^{\infty}\frac{H_n}{n^5}(-1)^{n-1}\end{aligned}$$

$$-5\sum_{n=1}^{\infty}\frac{L_n(1)}{n^5}(-1)^{n-1}-2\sum_{n=1}^{\infty}\frac{L_n(2)}{n^4}(-1)^{n-1}. \tag{3.39}$$

Noting that, using the following integrals

$$\int_0^1\frac{x^n\ln^m x}{1-x}dx=(-1)^m m!\left(\zeta(m+1)-\zeta_n(m+1)\right),\ 0<m\in\mathbb{Z},$$

$$\int_0^1\frac{x^n\ln^m x}{1+x}dx=(-1)^{m+n} m!\left(\bar{\zeta}(m+1)-L_n(m+1)\right),\ 0<m\in\mathbb{Z},$$

we can find that

$$\begin{aligned}\int_0^1\frac{\ln^m x\ln^2(1-x)}{1+x}dx=&2(-1)^m m!\left\{\sum_{n=1}^{\infty}\frac{H_n L_n(m+1)}{n}(-1)^{n-1}-\sum_{n=1}^{\infty}\frac{L_n(m+1)}{n^2}(-1)^{n-1}\right\}\\&+(-1)^m m!\ln^2 2\bar{\zeta}(m+1).\end{aligned} \tag{3.40}$$

$$\begin{aligned}\int_0^1\frac{\ln^m x\ln^2(1+x)}{1-x}dx=&2(-1)^m m!\left\{\sum_{n=1}^{\infty}\frac{H_n\zeta_n(m+1)}{n}(-1)^{n-1}-\sum_{n=1}^{\infty}\frac{\zeta_n(m+1)}{n^2}(-1)^{n-1}\right\}\\&+(-1)^m m!\ln^2 2\zeta(m+1).\end{aligned} \tag{3.41}$$

Therefore we immediately get the following Theorem.

Theorem 3.9 *For integer $m\geq 1$, then the integrals*

$$\int_0^1\frac{\ln^{2m}x\ln^2(1-x)}{1+x}dx,\int_0^1\frac{\ln^{2m}x\ln^2(1+x)}{1-x}dx$$

are expressible in terms of zeta values and linear sums.

Specially, taking $m=1$ in (3.40) (3.41) and combining (3.32) (3.34), we obtain

$$\int_0^1\frac{\ln^2 x\ln^2(1-x)}{1+x}dx=8\mathrm{Li}_5\left(\frac{1}{2}\right)-\frac{1}{15}\ln^5 2+\frac{15}{2}\zeta(5)+\frac{2}{3}\zeta(2)\ln^3 2-\frac{13}{2}\zeta(2)\zeta(3)+\zeta(4)\ln 2, \tag{3.42}$$

$$\begin{aligned}\int_0^1\frac{\ln^2 x\ln^2(1+x)}{1-x}dx=&-16\mathrm{Li}_5\left(\frac{1}{2}\right)-8\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)-\frac{1}{5}\ln^5 2+\frac{7}{2}\zeta(3)\ln^2 2+\frac{47}{2}\zeta(5)\\&+\frac{2}{3}\zeta(2)\ln^3 2-\frac{13}{4}\zeta(2)\zeta(3)-\frac{49}{4}\zeta(4)\ln 2.\end{aligned} \tag{3.43}$$

On the other hand, constructing function $y=\sum_{n=1}^{\infty}L_n^2(1)x^n\ (-1<x<1)$, we can easily obtain that

$$y=xy+2\sum_{n=1}^{\infty}\frac{L_n(1)(-1)^n}{n+1}x^{n+1}+\sum_{n=1}^{\infty}\frac{x^n}{n^2}. \tag{3.44}$$

Using the following integral

$$\int_0^x \frac{\ln(1-t)}{1+t}dt = \sum_{n=1}^{\infty} \frac{L_n(1)(-1)^n}{n+1}x^{n+1}.$$

Then we may rewrite (3.44) as

$$\sum_{n=1}^{\infty} \frac{\zeta_n(2) - L_n^2(1)}{n+1}x^{n+1} = 2\ln(1-x)\int_0^x \frac{\ln(1-t)}{1+t}dt - 2\int_0^x \frac{\ln^2(1-t)}{1+t}dt. \tag{3.45}$$

By using (2.11), the following identity is easily derived

$$\int_0^x \frac{\ln^2(1-t)}{1+t}dt = 2\sum_{n=1}^{\infty} \frac{H_n}{n+1}(-1)^{n+1}\left(\sum_{k=1}^{n+1}\frac{(-1)^k}{k}x^k + \ln(1+x)\right), x \in (-1,1]. \tag{3.46}$$

Substituting (3.46) into (3.45) respectively, we obtain

$$\begin{aligned}&\frac{1}{2}\sum_{n=1}^{\infty} \frac{L_n^2(1) - \zeta_n(2)}{n+1}x^{n+1} - \ln(1-x)\sum_{n=1}^{\infty} \frac{L_n(1)(-1)^{n+1}}{n+1}x^{n+1}\\ &= 2\sum_{n=1}^{\infty} \frac{H_n}{n+1}(-1)^{n+1}\left(\sum_{k=1}^{n+1}\frac{(-1)^k}{k}x^k + \ln(1+x)\right).\end{aligned} \tag{3.47}$$

Multiplying (3.47) by $\dfrac{\ln x}{x}$ and integrating over the interval (0,1). The result is

$$\begin{aligned}\sum_{n=1}^{\infty} \frac{L_n(1)\zeta_n(2)}{n^2}(-1)^{n-1} &= \frac{5}{4}\zeta(5) + \frac{15}{4}\zeta(4)\ln 2 - \frac{9}{8}\zeta(2)\zeta(3) + \frac{1}{2}\sum_{n=1}^{\infty} \frac{L_n^2(1)}{(n+1)^3}\\ &\quad - \sum_{n=1}^{\infty} \frac{H_n L_n(1)}{n^3}(-1)^{n-1} - \frac{1}{2}\int_0^1 \frac{\ln^2 x \ln^2(1-x)}{1+x}dx.\end{aligned} \tag{3.48}$$

Combining (3.30) (3.33) (3.42) with (3.48), we have

$$\begin{aligned}\sum_{n=1}^{\infty} \frac{L_n(1)\zeta_n(2)}{n^2}(-1)^{n-1} &= -4\mathrm{Li}_5\left(\frac{1}{2}\right) + \frac{1}{30}\ln^5 2 + \frac{7}{4}\zeta(3)\ln^2 2 - \frac{29}{64}\zeta(5)\\ &\quad - \frac{1}{3}\zeta(2)\ln^3 2 + \frac{15}{8}\zeta(2)\zeta(3) - \frac{23}{16}\zeta(4)\ln 2.\end{aligned} \tag{3.49}$$

In the same manner, replacing x by $-x$ in (3.47), then dividing this by $\dfrac{\ln x}{x}$ and integrating over the interval $(0,1)$, we arrive at the conclusion that

$$\begin{aligned}\sum_{n=1}^{\infty} \frac{L_n(1)L_n(2)}{n^2}(-1)^{n-1} &= 8\mathrm{Li}_5\left(\frac{1}{2}\right) + 4\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right) + \frac{1}{10}\ln^5 2 - \frac{35}{4}\zeta(5)\\ &\quad - \frac{1}{3}\zeta(2)\ln^3 2 + \frac{1}{16}\zeta(2)\zeta(3) + \frac{49}{8}\zeta(4)\ln 2.\end{aligned} \tag{3.50}$$

Applying (3.49) to (3.18), we obtain

$$\sum_{n=1}^{\infty}\frac{H_nL_n(2)}{n^2}=4\mathrm{Li}_5\left(\frac{1}{2}\right)+4\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)+\frac{2}{15}\ln^5 2+\frac{29}{64}\zeta(5)$$
$$+\frac{7}{4}\zeta(3)\ln^2 2-\frac{2}{3}\zeta(2)\ln^3 2-\frac{5}{4}\zeta(2)\zeta(3). \tag{3.51}$$

In many other cases we are also able to obtain a formula for other 5th order Euler sum explicitly in terms of values of zeta, logarithm and polylogarithm functions (See Table 5).

4 Triple Euler sums

We now define the generalized nth double harmonic number by

$$S_{r,p}(n)=\sum_{k=1}^{n}\frac{1}{k^p}\sum_{j=1}^{k}\frac{1}{j^r}=\sum_{k=1}^{n}\frac{\zeta_k(r)}{k^p}.$$

It is easy to see that, when $p>1$ is a positive integer

$$\lim_{n\to\infty}S_{r,p}(n)=S_{r,p}=\sum_{n=1}^{\infty}\frac{\zeta_n(r)}{n^p}.$$

For positive integers r,p,q, the linear triple Euler sums are defined by

$$S_{r,p,q}=\sum_{n=1}^{\infty}\frac{S_{r,p}(n)}{n^q}=\sum_{n=1}^{\infty}\frac{1}{n^q}\left(\sum_{k=1}^{n}\frac{\zeta_k(r)}{k^p}\right),q\geq 2.$$

By the definition of multiple zeta function and multiple zeta-star function, we have the relations

$$S_{r,p,q}=\zeta^{\star}(q,p,r),q\geq 2,$$

$$\zeta^{\star}(q,p,r)=\zeta(q,p,r)+\zeta^{\star}(q,p+r)+\zeta^{\star}(q+p,r)-\zeta(p+q+r). \tag{4.1}$$

The multiple zeta values $\zeta(q,p,r)$ are investigated [12, 19, 22]. Markett ([22]) gave explicit reductions to zeta values for all triple sums $\zeta(q,p,r)$ with $p+q+r\leq 6$, and he proved an explicit formula for $\zeta(q,1,1)$ in terms of zeta values. The best results to date are due to J. M. Borwein and R. Girgensohn ([12]). J.M. Borwein, R. Girgensohn proved that all $\zeta(q,p,r)$ with $r+p+q$ is even or less than or equal to 10 or $r+p+q=12$ were reducible to zeta values and linear double sums. By the relation (4.1), we have the following Theorem.

Theorem 4.1 *If $p+q+r$ is even or less than or equal to 10 or $p+q+r=12$, then triple linear Euler sums $S_{r,p,q}$ (or $\zeta^{\star}(q,p,r)$) can be expressed as a rational linear combination of products of zeta values and linear double Euler sums.*

Here are a few explicit evaluations.

$$S_{1,2,2}=\sum_{n=1}^{\infty}\frac{1}{n^2}\left(\sum_{k=1}^{n}\frac{H_k}{k^2}\right)=2\zeta(5),$$

Euler sums	Closed form
$\sum_{n=1}^{\infty}\frac{H_nL_n(1)}{n^3}$	$=2\mathrm{Li}_5\left(\frac{1}{2}\right)-\frac{1}{60}\ln^5 2-\frac{193}{64}\zeta(5)-\frac{7}{8}\zeta(3)\ln^2 2$ $+\frac{1}{6}\zeta(2)\ln^3 2+4\zeta(4)\ln 2+\frac{3}{8}\zeta(2)\zeta(3)$
$\sum_{n=1}^{\infty}\frac{L_n(1)\zeta_n(3)}{n}(-1)^{n-1}$	$=-2\mathrm{Li}_5\left(\frac{1}{2}\right)+\frac{1}{60}\ln^5 2-\frac{3}{8}\zeta(3)\ln^2 2-\frac{193}{64}\zeta(5)$ $-\frac{1}{6}\zeta(2)\ln^3 2+\frac{11}{16}\zeta(4)\ln 2-\frac{5}{8}\zeta(2)\zeta(3)$
$\sum_{n=1}^{\infty}\frac{L_n(1)L_n(3)}{n}(-1)^{n-1}$	$=2\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)+\frac{1}{12}\ln^5 2+\frac{3}{8}\zeta(3)\ln^2 2-\frac{19}{32}\zeta(5)$ $-\frac{1}{2}\zeta(2)\ln^3 2+\frac{11}{16}\zeta(4)\ln 2+\frac{1}{4}\zeta(2)\zeta(3)$
$\sum_{n=1}^{\infty}\frac{L_n^4(1)}{n+1}(-1)^{n-1}$	$=\frac{11}{30}\ln^5 2-4\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)-\frac{73}{2}\zeta(4)\ln 2+7\zeta(2)\ln^3 2$ $+\frac{5}{8}\zeta(2)\zeta(3)+\frac{709}{16}\zeta(5)-48\mathrm{Li}_5\left(\frac{1}{2}\right)$
$\sum_{n=1}^{\infty}\frac{L_n(1)\zeta_n(2)}{n^2}$	$=-8\mathrm{Li}_5\left(\frac{1}{2}\right)-4\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)-\frac{1}{10}\ln^5 2+\frac{27}{4}\zeta(5)$ $+\frac{1}{3}\zeta(2)\ln^3 2+\frac{1}{2}\zeta(2)\zeta(3)-\frac{23}{16}\zeta(4)\ln 2$
$\sum_{n=1}^{\infty}\frac{\zeta_n^2(2)}{n}(-1)^{n-1}$	$=16\mathrm{Li}_5\left(\frac{1}{2}\right)+8\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)+\frac{1}{5}\ln^5 2-\frac{259}{16}\zeta(5)$ $-\frac{2}{3}\zeta(2)\ln^3 2+\frac{5}{8}\zeta(2)\zeta(3)+\frac{29}{4}\zeta(4)\ln 2$
$\sum_{n=1}^{\infty}\frac{H_n^4}{n}(-1)^{n-1}$	$=8\mathrm{Li}_5\left(\frac{1}{2}\right)+4\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)+\frac{3}{10}\ln^5 2+\frac{9}{4}\zeta(3)\ln^2 2$ $-\frac{83}{16}\zeta(5)-\frac{4}{3}\zeta(2)\ln^3 2-\frac{11}{8}\zeta(2)\zeta(3)+\frac{11}{4}\zeta(4)\ln 2$
$\sum_{n=1}^{\infty}\frac{H_n^2\zeta_n(2)}{n}(-1)^{n-1}$	$=-2\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)-\frac{1}{12}\ln^5 2-\frac{7}{8}\zeta(3)\ln^2 2+\frac{1}{3}\zeta(2)\ln^3 2$ $+\frac{3}{8}\zeta(2)\zeta(3)+\frac{7}{8}\zeta(4)\ln 2$
$\sum_{n=1}^{\infty}\frac{H_n^2L_n(1)}{n^2}$	$=4\mathrm{Li}_5\left(\frac{1}{2}\right)+2\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)+\frac{1}{20}\ln^5 2-\frac{165}{32}\zeta(5)-\frac{1}{6}\zeta(2)\ln^3 2$ $+\frac{1}{2}\zeta(2)\zeta(3)+\frac{109}{16}\zeta(4)\ln 2$
$\sum_{n=1}^{\infty}\frac{L_n^2(2)}{n}(-1)^{n-1}$	$=-16\mathrm{Li}_5\left(\frac{1}{2}\right)-8\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)-\frac{1}{5}\ln^5 2+\frac{299}{16}\zeta(5)$ $+\frac{2}{3}\zeta(2)\ln^3 2-\frac{1}{4}\zeta(2)\zeta(3)-\frac{39}{4}\zeta(4)\ln 2$
$\sum_{n=1}^{\infty}\frac{H_n^2L_n(1)}{n^2}(-1)^{n-1}$	$=2\mathrm{Li}_5\left(\frac{1}{2}\right)-\frac{1}{60}\ln^5 2-\frac{7}{8}\zeta(3)\ln^2 2-\frac{197}{64}\zeta(5)$ $+\frac{1}{6}\zeta(2)\ln^3 2-\frac{3}{4}\zeta(2)\zeta(3)+\frac{109}{16}\zeta(4)\ln 2$
$\sum_{n=1}^{\infty}\frac{H_nL_n(2)}{n^2}(-1)^{n-1}$	$=\frac{125}{32}\zeta(5)-\frac{13}{8}\zeta(2)\zeta(3)$

$$\begin{array}{ll}
\sum_{n=1}^{\infty}\frac{L_n^2(1)L_n(2)}{n}(-1)^{n-1} & =14\mathrm{Li}_5\left(\frac{1}{2}\right)-\frac{7}{60}\ln^5 2-\frac{7}{8}\zeta(3)\ln^2 2-\frac{899}{64}\zeta(5)-\frac{2}{3}\zeta(2)\ln^3 2 \\
& \quad +\frac{1}{4}\zeta(2)\zeta(3)+\frac{193}{16}\zeta(4)\ln 2 \\
\sum_{n=1}^{\infty}\frac{L_n^2(1)\zeta_n(2)}{n}(-1)^{n-1} & =-6\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)-\frac{1}{4}\ln^5 2-\frac{1}{6}\zeta(2)\ln^3 2+\frac{1}{8}\zeta(2)\zeta(3) \\
& \quad +\frac{9}{2}\zeta(4)\ln 2 \\
\sum_{n=1}^{\infty}\frac{\zeta_n(2)L_n(2)}{n}(-1)^{n-1} & =\frac{75}{64}\zeta(5)+\frac{3}{8}\zeta(2)\zeta(3)-\frac{25}{16}\zeta(4)\ln 2 \\
\sum_{n=1}^{\infty}\frac{H_n^3 L_n(1)}{n}(-1)^{n-1} & =-2\mathrm{Li}_5\left(\frac{1}{2}\right)-\frac{11}{60}\ln^5 2-\frac{9}{16}\zeta(3)\ln^2 2+\frac{55}{16}\zeta(5)+\frac{7}{12}\zeta(2)\ln^3 2 \\
& \quad -\frac{1}{16}\zeta(2)\zeta(3)+\frac{37}{8}\zeta(4)\ln 2 \\
\sum_{n=1}^{\infty}\frac{H_n L_n(1)\zeta_n(2)}{n}(-1)^{n-1} & =-6\mathrm{Li}_5\left(\frac{1}{2}\right)+\frac{1}{20}\ln^5 2+\frac{7}{16}\zeta(3)\ln^2 2+\frac{155}{32}\zeta(5) \\
& \quad -\frac{1}{12}\zeta(2)\ln^3 2+\frac{7}{16}\zeta(2)\zeta(3)-2\zeta(4)\ln 2
\end{array}$$

TABLE5. Some closed form of nonlinear Euler sums

$$\begin{aligned}
S_{2,1,2} &= \sum_{n=1}^{\infty}\frac{1}{n^2}\left(\sum_{k=1}^{n}\frac{\zeta_k(2)}{k}\right)=\frac{9}{2}\zeta(5)-\zeta(2)\zeta(3), \\
S_{1,2,3} &= \sum_{n=1}^{\infty}\frac{1}{n^3}\left(\sum_{k=1}^{n}\frac{H_k}{k^2}\right)=3\zeta^2(3)-\frac{143}{48}\zeta(6), \\
S_{2,1,3} &= \sum_{n=1}^{\infty}\frac{1}{n^3}\left(\sum_{k=1}^{n}\frac{\zeta_k(2)}{k}\right)=\frac{11}{8}\zeta(6), \\
S_{1,3,2} &= \sum_{n=1}^{\infty}\frac{1}{n^2}\left(\sum_{k=1}^{n}\frac{H_k}{k^3}\right)=\frac{145}{24}\zeta(6)-3\zeta^2(3), \\
S_{1,2,4} &= \sum_{n=1}^{\infty}\frac{1}{n^4}\left(\sum_{k=1}^{n}\frac{H_k}{k^2}\right)=\frac{115}{16}\zeta(7)+\frac{5}{2}\zeta(3)\zeta(4)-\frac{11}{2}\zeta(2)\zeta(5), \\
S_{2,1,4} &= \sum_{n=1}^{\infty}\frac{1}{n^4}\left(\sum_{k=1}^{n}\frac{\zeta_k(2)}{k}\right)=\frac{61}{8}\zeta(7)-\frac{5}{2}\zeta(2)\zeta(5)-\frac{7}{4}\zeta(3)\zeta(4).
\end{aligned}$$

Next, we evaluate some specific nonlinear triple Euler sums involving harmonic numbers and double harmonic numbers.

Theorem 4.2 *For $p,q,r>0$, then*

$$S_{r,p}(n)\zeta_n(q)+S_{r,p+q}(n)=\sum_{k=1}^{n}\frac{S_{r,p}(k)}{k^q}+\sum_{k=1}^{n}\frac{\zeta_k(r)\zeta_k(q)}{k^p}. \tag{4.2}$$

Proof. Constructing function

$$y=\sum_{n=1}^{\infty}S_{r,p}\left(n\right)\zeta_{n}\left(q\right)x^{n-1}\quad\left(\left|x\right|<1\right). \tag{4.3}$$

By using the definition of the harmonic numbers $S_{r,p}(n)$ and $\zeta_n(q)$, we have the relations

$$S_{r,p}\left(n+1\right)=S_{r,p}\left(n\right)+\frac{\zeta_{n+1}\left(r\right)}{\left(n+1\right)^{p}},\zeta_{n+1}\left(q\right)=\zeta_{n}\left(q\right)+\frac{1}{\left(n+1\right)^{q}}. \tag{4.4}$$

Applying (4.4), we may rewrite (4.3) as

$$y=xy+\sum_{n=1}^{\infty}\left\{\frac{S_{r,p}\left(n\right)}{n^{q}}-\frac{\zeta_{n}\left(r\right)}{n^{p+q}}+\frac{\zeta_{n}\left(r\right)\zeta_{n}\left(q\right)}{n^{p}}\right\}x^{n-1}, \tag{4.5}$$

which, moved xy from the right to the left and multiplied it by $(1-x)^{-1}$. Using Cauchy product formula, we get

$$\sum_{n=1}^{\infty}S_{r,p}\left(n\right)\zeta_{n}\left(q\right)x^{n-1}=\sum_{n=1}^{\infty}\sum_{k=1}^{n}\left\{\frac{S_{r,p}\left(k\right)}{k^{q}}-\frac{\zeta_{k}\left(r\right)}{k^{p+q}}+\frac{\zeta_{k}\left(r\right)\zeta_{k}\left(q\right)}{k^{p}}\right\}x^{n-1}. \tag{4.6}$$

Equating coefficients of x^{n-1} in (4.6), we can obtain (4.2). □

More generally, constructing function

$$y=\sum_{n=1}^{\infty}\left\{\zeta_{n}\left(m\right)\left(\sum_{k=1}^{n}\frac{\prod_{i=1}^{t}\zeta_{k}\left(p_{i}\right)}{k^{s}}\right)-\left(\sum_{k=1}^{n}\frac{\prod_{i=1}^{t}\zeta_{k}\left(p_{i}\right)}{k^{s+m}}\right)\right\}x^{n-1},\ \left|x\right|<1,$$

we have

$$\sum_{n=1}^{\infty}\frac{1}{n^{m}}\left(\sum_{k=1}^{n}\frac{\prod_{i=1}^{t}\zeta_{k}\left(p_{i}\right)}{k^{s}}\right)+\sum_{n=1}^{\infty}\frac{\zeta_{n}\left(m\right)\prod_{i=1}^{t}\zeta_{n}\left(p_{i}\right)}{n^{s}}=\sum_{n=1}^{\infty}\frac{\prod_{i=1}^{t}\zeta_{n}\left(p_{i}\right)}{n^{s+m}}+\zeta\left(m\right)\sum_{n=1}^{\infty}\frac{\prod_{i=1}^{t}\zeta_{n}\left(p_{i}\right)}{n^{s}}, \tag{4.7}$$

where $m,s>1$. Taking $m=s=t=2,p_i=1,i=1,2$ in (4.7), we obtain

$$\sum_{n=1}^{\infty}\frac{1}{n^{2}}\left(\sum_{k=1}^{n}\frac{H_{k}^{2}}{k^{2}}\right)=\frac{129}{16}\zeta\left(6\right)-4\zeta^{2}\left(3\right).$$

Let $t=2,m=s=p_1=p_2=l>1$ in (4.7), we get

$$\sum_{n=1}^{\infty}\frac{1}{n^{l}}\left(\sum_{k=1}^{n}\frac{\zeta_{k}^{2}\left(l\right)}{k^{l}}\right)=\sum_{n=1}^{\infty}\frac{\zeta_{n}^{2}\left(l\right)}{n^{2l}}+\zeta\left(l\right)\sum_{n=1}^{\infty}\frac{\zeta_{n}^{2}\left(l\right)}{n^{l}}-\sum_{n=1}^{\infty}\frac{\zeta_{n}^{3}\left(l\right)}{n^{l}}.$$

From [19, 34], we know that the quadratic sums $\sum_{n=1}^{\infty}\frac{\zeta_{n}^{2}\left(l\right)}{n^{2l}}$ and the cubic sums $S_{l^3,l}=\sum_{n=1}^{\infty}\frac{\zeta_{n}^{3}\left(l\right)}{n^{l}}$ are reducible to linear sums. Hence, we obtain the following Theorem.

Theorem 4.3 *For integer $l>1$, the triple sums*

$$\sum_{n=1}^{\infty}\frac{1}{n^{l}}\left(\sum_{k=1}^{n}\frac{\zeta_{k}^{2}\left(l\right)}{k^{l}}\right)$$

are reducible to linear double sums.

As an example, we hvae

$$\sum_{n=1}^{\infty}\frac{1}{n^{2}}\left(\sum_{k=1}^{n}\frac{\zeta_{k}^{2}\left(2\right)}{k^{2}}\right)=-\frac{9}{2}S_{2,6}-\frac{1291}{96}\zeta\left(8\right)-2\zeta\left(2\right)\zeta^{2}\left(3\right)+20\zeta\left(3\right)\zeta\left(5\right).$$

Similarly, constructing function

$$y=\sum_{n=1}^{\infty}S_{r,p}\left(n\right)S_{l,m}\left(n\right)x^{n-1},y=\sum_{n=1}^{\infty}S_{r,p}\left(n\right)\zeta_{n}^{2}\left(q\right)x^{n-1}\quad\left(\left|x\right|<1\right),$$

we can obtain the following Theorems.

Theorem 4.4 *For $p,r,m,l>0$, we have*

$$S_{r,p}\left(n\right)S_{l,m}\left(n\right)+\sum_{k=1}^{n}\frac{\zeta_{k}\left(r\right)\zeta_{k}\left(l\right)}{k^{p+m}}=\sum_{k=1}^{n}\frac{S_{r,p}\left(k\right)\zeta_{k}\left(l\right)}{k^{m}}+\sum_{k=1}^{n}\frac{S_{l,m}\left(k\right)\zeta_{k}\left(r\right)}{k^{p}}. \tag{4.8}$$

Theorem 4.5 *For $p,r,q>0$, we have*

$$\begin{aligned}&S_{r,p}\left(n\right)\zeta_{n}^{2}\left(q\right)+2\sum_{k=1}^{n}\frac{\zeta_{k}\left(r\right)\zeta_{k}\left(q\right)}{k^{p+q}}+\sum_{k=1}^{n}\frac{S_{r,p}\left(k\right)}{k^{2q}}\\&=2\sum_{k=1}^{n}\frac{S_{r,p}\left(k\right)\zeta_{k}\left(q\right)}{k^{q}}+\sum_{k=1}^{n}\frac{\zeta_{k}^{2}\left(q\right)\zeta_{k}\left(r\right)}{k^{p}}+S_{r,p+2q}\left(n\right).\end{aligned} \tag{4.9}$$

Letting $n\to\infty$ in (4.2) (4.8) and (4.9), we obtain that for $p>1,q>1,m>1,l>0,r>0$,

$$S_{r,p}\zeta\left(q\right)+S_{r,p+q}=S_{r,p,q}+S_{rq,p}, \tag{4.10}$$

$$S_{r,p}S_{l,m}+S_{rl,p+m}=\sum_{n=1}^{\infty}\frac{S_{r,p}\left(n\right)\zeta_{n}\left(l\right)}{n^{m}}+\sum_{n=1}^{\infty}\frac{S_{l,m}\left(n\right)\zeta_{n}\left(r\right)}{n^{p}}, \tag{4.11}$$

$$S_{r,p}\zeta^{2}\left(q\right)+2S_{rq,p+q}+S_{r,p,2q}=2\sum_{n=1}^{\infty}\frac{S_{r,p}\left(n\right)\zeta_{n}\left(q\right)}{n^{q}}+S_{q^{2}r,p}+S_{r,p+2q}, \tag{4.12}$$

Here $S_{q^{2}r,p}=\sum_{k=1}^{n}\frac{\zeta_{k}^{2}\left(q\right)\zeta_{k}\left(r\right)}{k^{p}}$, $S_{rl,p+m}=\sum_{n=1}^{\infty}\frac{\zeta_{n}\left(r\right)\zeta_{n}\left(l\right)}{n^{p+m}}$, $S_{rq,p+q}=\sum_{n=1}^{\infty}\frac{\zeta_{n}\left(r\right)\zeta_{n}\left(q\right)}{n^{p+q}}$.
From Theorem 2.8 in the reference [34], we deduce that

$$\sum_{k=1}^{n}\frac{H_{k}^{2}}{k}=\frac{1}{3}H_{n}^{3}+\sum_{k=1}^{n}\frac{H_{k}}{k^{2}}-\frac{1}{3}\zeta_{n}\left(3\right), \tag{4.13}$$

$$H_{n}^{4}=4\sum_{k=1}^{n}\frac{H_{k}^{3}}{k}-6\sum_{k=1}^{n}\frac{H_{k}^{2}}{k^{2}}+4\sum_{k=1}^{n}\frac{H_{k}}{k^{3}}-\zeta_{n}\left(4\right), \tag{4.14}$$

$$H_n^2\zeta_n(2)=\sum_{k=1}^{n}\frac{H_k^2}{k^2}+2\sum_{k=1}^{n}\frac{H_k\zeta_k(2)}{k}-2\sum_{k=1}^{n}\frac{H_k}{k^3}-\sum_{k=1}^{n}\frac{\zeta_k(2)}{k^2}+\zeta_n(4). \tag{4.15}$$

Multiplying (4.13)-(4.15) by $\frac{1}{n^m}$ $(m>1)$ and summing with respect to n, the result is

$$\sum_{n=1}^{\infty}\frac{1}{n^m}\left(\sum_{k=1}^{n}\frac{H_k^2}{k}\right)=\frac{1}{3}S_{1^3,m}+S_{1,2,m}-\frac{1}{3}S_{3,m}, \tag{4.16}$$

$$\sum_{n=1}^{\infty}\frac{1}{n^m}\left(\sum_{k=1}^{n}\frac{H_k^3}{k}\right)-\frac{3}{2}\sum_{n=1}^{\infty}\frac{1}{n^m}\left(\sum_{k=1}^{n}\frac{H_k^2}{k^2}\right)=\frac{1}{4}S_{1^4m}-S_{1,3,m}+\frac{1}{4}S_{4,m}, \tag{4.17}$$

$$\sum_{n=1}^{\infty}\frac{1}{n^m}\left(\sum_{k=1}^{n}\frac{H_k\zeta_k(2)}{k}\right)+\frac{1}{2}\sum_{n=1}^{\infty}\frac{1}{n^m}\left(\sum_{k=1}^{n}\frac{H_k^2}{k^2}\right)=\frac{1}{2}S_{1^2,2,m}+S_{1,3,m}+\frac{1}{2}S_{2,2,m}-\frac{1}{2}S_{4,m}. \tag{4.18}$$

Combining related equation, we conclude that

$$\begin{aligned}
&\sum_{n=1}^{\infty}\frac{1}{n^2}\left(\sum_{k=1}^{n}\frac{H_k^2}{k}\right)=\frac{7}{2}\zeta(5)+\zeta(2)\zeta(3),\\
&\sum_{n=1}^{\infty}\frac{1}{n^3}\left(\sum_{k=1}^{n}\frac{H_k^2}{k}\right)=2\zeta^2(3)-\frac{29}{24}\zeta(6),\\
&\sum_{n=1}^{\infty}\frac{1}{n^4}\left(\sum_{k=1}^{n}\frac{H_k^2}{k}\right)=6\zeta(7)-\frac{3}{2}\zeta(2)\zeta(5)-\frac{7}{4}\zeta(3)\zeta(4),\\
&\sum_{n=1}^{\infty}\frac{1}{n^2}\left(\sum_{k=1}^{n}\frac{H_k^3}{k}\right)=\frac{1631}{96}\zeta(6)-\frac{5}{2}\zeta^2(3),\\
&\sum_{n=1}^{\infty}\frac{1}{n^2}\left(\sum_{k=1}^{n}\frac{H_k\zeta_k(2)}{k}\right)=\frac{151}{48}\zeta(6)+\frac{1}{2}\zeta^2(3).
\end{aligned}$$

Taking $r=1,p=m=l=2$ in (4.11), we have

$$\begin{aligned}
\sum_{n=1}^{\infty}\frac{\zeta_n(2)}{n^2}\left(\sum_{k=1}^{n}\frac{H_k}{k^2}\right)=&\left(\sum_{n=1}^{\infty}\frac{H_n}{n^2}\right)\left(\sum_{n=1}^{\infty}\frac{\zeta_n(2)}{n^2}\right)+\sum_{n=1}^{\infty}\frac{H_n\zeta_n(2)}{n^4}\\
&-\frac{1}{2}\sum_{n=1}^{\infty}\frac{H_n\zeta_n^2(2)}{n^2}-\frac{1}{2}\sum_{n=1}^{\infty}\frac{H_n\zeta_n(4)}{n^2}\\
=&\frac{83}{16}\zeta(7)-\frac{7}{2}\zeta(2)\zeta(5)+\frac{5}{2}\zeta(3)\zeta(4).
\end{aligned}$$

Similarly, we can obtain that

$$\begin{aligned}
&\sum_{n=1}^{\infty}\frac{\zeta_n(2)}{n^2}\left(\sum_{k=1}^{n}\frac{H_k^2}{k}\right)=\frac{67}{16}\zeta(7)+\frac{15}{2}\zeta(3)\zeta(4)-\frac{7}{2}\zeta(2)\zeta(5),\\
&\sum_{n=1}^{\infty}\frac{\zeta_n(2)}{n^2}\left(\sum_{k=1}^{n}\frac{\zeta_k(2)}{k}\right)=-\frac{169}{16}\zeta(7)+\frac{15}{2}\zeta(2)\zeta(5)+\zeta(3)\zeta(4).
\end{aligned}$$

For triple alternating Euler sums, by constructing function, we also give the following results.

$$\sum_{n=1}^{\infty}\frac{(-1)^{n-1}}{n^2}\left(\sum_{k=1}^{n}\frac{L_k(1)}{k}\right)=\frac{3}{16}\zeta(4)+\frac{3}{4}\zeta(2)\ln^2 2,$$

$$\sum_{n=1}^{\infty}\frac{(-1)^{n-1}}{n^2}\left(\sum_{k=1}^{n}\frac{H_k}{k}(-1)^{k-1}\right)=\frac{39}{16}\zeta(4)-\frac{1}{8}\ln^4 2-3\mathrm{Li}_4\left(\frac{1}{2}\right),$$

$$\sum_{n=1}^{\infty}\frac{1}{n^2}\left(\sum_{k=1}^{n}\frac{L_k(1)}{k}\right)=3\zeta(4)+\frac{3}{2}\zeta(2)\ln^2 2-\frac{1}{8}\ln^4 2-\frac{7}{8}\zeta(3)\ln 2-3\mathrm{Li}_4\left(\frac{1}{2}\right),$$

$$\sum_{n=1}^{\infty}\frac{1}{n^2}\left(\sum_{k=1}^{n}\frac{H_k}{k}(-1)^{k-1}\right)=5\zeta(4)+\frac{1}{4}\zeta(2)\ln^2 2-\frac{1}{6}\ln^4 2-\frac{21}{8}\zeta(3)\ln 2-4\mathrm{Li}_4\left(\frac{1}{2}\right),$$

$$\begin{aligned}\sum_{n=1}^{\infty}\frac{(-1)^{n-1}}{n^3}\left(\sum_{k=1}^{n}\frac{H_k}{k}(-1)^{k-1}\right)&=-2\mathrm{Li}_5\left(\frac{1}{2}\right)+\frac{1}{60}\ln^5 2+\frac{85}{16}\zeta(5)-\frac{1}{6}\zeta(2)\ln^3 2\\&\quad-\frac{5}{4}\zeta(2)\zeta(3)-\frac{1}{4}\zeta(4)\ln 2\end{aligned}$$

$$\begin{aligned}\sum_{n=1}^{\infty}\frac{1}{n^3}\left(\sum_{k=1}^{n}\frac{H_k}{k}(-1)^{k-1}\right)&=4\mathrm{Li}_5\left(\frac{1}{2}\right)+2\ln 2\mathrm{Li}_4\left(\frac{1}{2}\right)+\frac{1}{20}\ln^5 2-\frac{7}{8}\zeta(3)\ln^2 2-\frac{27}{8}\zeta(5)\\&\quad+\frac{1}{6}\zeta(2)\ln^3 2-\frac{1}{16}\zeta(2)\zeta(3)-\frac{49}{16}\zeta(4)\ln 2.\end{aligned}$$

5 Some evaluation of Euler-type sums

In this section, we will establish some explicit relationships which involve Euler sums of the form

$$S_{m_1m_2\ldots m_r}(r,k)=\sum_{n=1}^{\infty}\frac{\zeta_n(m_1)\cdots\zeta_n(m_r)}{(n+r)(n+k)},\ 1\le k,r\in\mathbb{Z}$$

and harmonic numbers. We develops an approach to evaluation of Euler-type sums $S_{m_1m_2\ldots m_r}(r,k)$. The approach is based on simple Cauchy product formula and integral computations. We can find that many Euler-type sums $S_{m_1m_2\ldots m_r}(r,k)$ can be expressed in terms of series of Riemann zeta function values and harmonic numbers. Now we state our main theorems.

Theorem 5.1 *For integers $m,k>0$, then*

$$\sum_{n=1}^{\infty}\frac{\zeta_n(m)}{n(n+k)}=\frac{1}{k}\left\{\zeta(m+1)+\sum_{j=1}^{m-1}(-1)^{j-1}\zeta(m+1-j)\zeta_{k-1}(j)+(-1)^{m-1}\sum_{i=1}^{k-1}\frac{H_i}{i^m}\right\}.\quad(5.1)$$

Proof. By the definition of polylogarithm function and Cauchy product formula, we can verify that

$$\frac{\mathrm{Li}_m(x)}{1-x}=\sum_{n=1}^{\infty}\zeta_n(m)x^n,\ x\in(-1,1).\quad(5.2)$$

Multiplying (5.2) by $x^{r-1}-x^{k-1}$ and integrating over (0,1), we obtain

$$\int_0^1\left(x^{r-1}-x^{k-1}\right)\frac{\mathrm{Li}_m(x)}{1-x}dx=(k-r)\sum_{n=1}^{\infty}\frac{\zeta_n(m)}{(n+r)(n+k)}\quad(0\le r<k,\ r,k\in Z).\quad(5.3)$$

We now evaluate the integral on the left side of (5.3). Noting that

$$\int_0^1 \left(x^{r-1}-x^{k-1}\right)\frac{\mathrm{Li}_m(x)}{1-x}dx=\sum_{i=1}^{k-r}\int_0^1 x^{r+i-2}\mathrm{Li}_m(x)\,dx. \tag{5.4}$$

Using integration by parts we have

$$\int_0^1 x^{n-1}\mathrm{Li}_q(x)dx=\sum_{i=1}^{q-1}\frac{(-1)^{i-1}}{n^i}\zeta(q+1-i)+\frac{(-1)^{q-1}}{n^q}H_n, 1\le q\in\mathbb{Z}. \tag{5.5}$$

Taking $r=0$ in (5.3) and (5.4), substituting (5.5) into (5.3) and (5.4), we can obtain (5.1). □
If taking $r\ge 1$ in (5.3) and (5.4), using (5.5), we have

$$\sum_{n=1}^{\infty}\frac{\zeta_n(m)}{(n+r)(n+k)}=\frac{1}{k-r}\left\{\begin{array}{l}\displaystyle\sum_{j=1}^{m-1}(-1)^{j-1}\zeta(m+1-j)\left(\zeta_{k-1}(j)-\zeta_{r-1}(j)\right)\\ \displaystyle+(-1)^{m-1}\left(\sum_{i=1}^{k-1}\frac{H_i}{i^m}-\sum_{i=1}^{r-1}\frac{H_i}{i^m}\right)\end{array}\right\}. \tag{5.6}$$

Let $m=1$ in (5.1), we can obtain the following formula

$$\sum_{n=1}^{\infty}\frac{H_n}{n(n+k)}=\frac{1}{k}\left(\frac{1}{2}H_k^2+\frac{1}{2}\zeta_k(2)+\zeta(2)-\frac{H_k}{k}\right). \tag{5.7}$$

Similarly to the proof of Theorem 5.1, we obtain the following formulae.

$$\sum_{n=1}^{\infty}\frac{L_n(m)}{n(n+k)}=\frac{1}{k}\left\{\begin{array}{l}\bar{\zeta}(m+1)+\displaystyle\sum_{j=1}^{m-1}(-1)^{j-1}\bar{\zeta}(m+1-j)\,\zeta_{k-1}(j)\\ \displaystyle+(-1)^{m-1}\ln 2\left(\zeta_{k-1}(m)+L_{k-1}(m)\right)+(-1)^m\sum_{i=1}^{k-1}\frac{(-1)^{i-1}}{i^m}L_i(1)\end{array}\right\}, \tag{5.8}$$

$$\sum_{n=1}^{\infty}\frac{L_n(m)}{(n+r)(n+k)}=\frac{1}{k-r}\left\{\begin{array}{l}\displaystyle\sum_{j=1}^{m-1}(-1)^{j-1}\bar{\zeta}(m+1-j)\left(\zeta_{k-1}(j)-\zeta_{r-1}(j)\right)\\ \displaystyle+(-1)^{m-1}\ln 2\left(\zeta_{k-1}(m)-\zeta_{r-1}(m)+L_{k-1}(m)-L_{r-1}(m)\right)\\ \displaystyle+(-1)^m\sum_{i=r}^{k-1}\frac{(-1)^{i-1}}{i^m}L_i(1)\end{array}\right\}. \tag{5.9}$$

Taking $m=1$ in (5.8), we have

$$\sum_{n=1}^{\infty}\frac{L_n(1)}{n(n+k)}=\frac{1}{k}\left\{\begin{array}{l}\bar{\zeta}(2)+\ln 2\left(H_k+L_k(1)\right)-\ln 2\dfrac{1+(-1)^{k-1}}{k}\\ \displaystyle-\frac{1}{2}\left(L_k^2(1)+\zeta_k(2)\right)+\frac{L_k(1)}{k}(-1)^{k-1}\end{array}\right\}. \tag{5.10}$$

Next, we consider the following quadratic Euler-type sums

$$\sum_{n=1}^{\infty}\frac{\zeta_n(m_1)\,\zeta_n(m_2)}{(n+r)(n+k)},\ 0\le r<k\in\mathbb{Z},\ 1\le m_i\in\mathbb{Z}.$$

Theorem 5.2 *For integers $m_1>0, m_2>0, k\ge 1, r\ge 0$, we have*

$$\begin{aligned}&(k-r)\sum_{n=1}^{\infty}\frac{\zeta_n(m_1)\,\zeta_n(m_2)-\zeta_n(m_1+m_2)}{(n+r)(n+k)}\\&=\sum_{j=1}^{k-r}\sum_{n=1}^{\infty}\left\{\frac{\zeta_n(m_1)}{n^{m_2}(n+r+j-1)}+\frac{\zeta_n(m_2)}{n^{m_2}(n+r+j-1)}-\frac{2}{n^{m_1+m_2}(n+r+j-1)}\right\}.\end{aligned}\tag{5.11}$$

where the Euler-type sums $\sum_{n=1}^{\infty}\frac{\zeta_n(m)}{n^p(n+j)}$ $(1\le p,m,j\in\mathbb{Z})$ can be evaluated by the following formula

$$\sum_{n=1}^{\infty}\frac{\zeta_n(m)}{n^p(n+j)}=\sum_{i=1}^{p-1}\frac{(-1)^{i-1}}{j^i}\sum_{n=1}^{\infty}\frac{\zeta_n(m)}{n^{p+1-i}}+\frac{(-1)^{p-1}}{j^{p-1}}\sum_{n=1}^{\infty}\frac{\zeta_n(m)}{n(n+j)}.\tag{5.12}$$

Proof. Similarly to the proof of Theorem 4.2, we construct function

$$y=\sum_{n=1}^{\infty}\left\{\zeta_n(m_1)\,\zeta_n(m_2)-\zeta_n(m_1+m_2)\right\}x^{n-1},\ |x|<1,$$

then have

$$\begin{aligned}&\sum_{n=1}^{\infty}\left\{\zeta_n(m_1)\,\zeta_n(m_2)-\zeta_n(m_1+m_2)\right\}x^{n-1}\\&=\sum_{n=1}^{\infty}\left\{\frac{\zeta_n(m_1)}{n^{m_2}}+\frac{\zeta_n(m_2)}{n^{m_1}}-\frac{2}{n^{m_1+m_2}}\right\}\frac{x^{n-1}}{1-x}.\end{aligned}\tag{5.13}$$

Multiplying (5.13) by x^r-x^k and integrating over (0,1), we can obtain (5.11). □
Let $r=0$ in(5.11), we have

$$\begin{aligned}&k\sum_{n=1}^{\infty}\frac{\zeta_n(m_1)\,\zeta_n(m_2)-\zeta_n(m_1+m_2)}{n(n+k)}\\&=\sum_{n=1}^{\infty}\left\{\frac{\zeta_n(m_1)}{n^{m_2+1}}+\frac{\zeta_n(m_2)}{n^{m_1+1}}-\frac{2}{n^{m_1+m_2+1}}\right\}\\&\quad+\sum_{j=1}^{k-1}\sum_{n=1}^{\infty}\left\{\frac{\zeta_n(m_1)}{n^{m_2}(n+j)}+\frac{\zeta_n(m_2)}{n^{m_2}(n+j)}-\frac{2}{n^{m_1+m_2}(n+j)}\right\},\end{aligned}\tag{5.14}$$

Proceeding in a similar fashion to evaluation of the Theorem 5.2, by constructing functions

$$y=\sum_{n=1}^{\infty}\left\{L_n(m_1)\,L_n(m_2)-\zeta_n(m_1+m_2)\right\}x^{n-1},\ |x|<1,$$

$$y=\sum_{n=1}^{\infty}\left\{\zeta_n(m_1)\,L_n(m_2)-L_n(m_1+m_2)\right\}x^{n-1},\ |x|<1,$$
$$y=\sum_{n=1}^{\infty}\left\{\zeta_n(m_1)\,\zeta_n(m_2)\,\zeta_n(m_3)-\zeta_n(m_1+m_2+m_3)\right\}x^{n-1},\ |x|<1,$$

we can obtain the following Theorems.

Theorem 5.3 *For integers $m_1>0,m_2>0,k\geq 1,r\geq 0$, we have*

$$\begin{aligned}&(k-r)\sum_{n=1}^{\infty}\frac{L_n(m_1)\,L_n(m_2)-\zeta_n(m_1+m_2)}{(n+r)(n+k)}\\&=\sum_{j=1}^{k-r}\sum_{n=1}^{\infty}\left\{\frac{L_n(m_1)}{n^{m_2}(n+r+j-1)}+\frac{L_n(m_2)}{n^{m_2}(n+r+j-1)}-\frac{2}{n^{m_1+m_2}(n+r+j-1)}\right\}.\end{aligned}$$

Theorem 5.4 *For integers $m_1>0,m_2>0,k\geq 1,r\geq 0$, we have*

$$\begin{aligned}&(k-r)\sum_{n=1}^{\infty}\frac{\zeta_n(m_1)\,L_n(m_2)-L_n(m_1+m_2)}{(n+r)(n+k)}\\&=\sum_{j=1}^{k-r}\sum_{n=1}^{\infty}\left\{\frac{\zeta_n(m_1)}{n^{m_2}(n+r+j-1)}(-1)^{n-1}+\frac{L_n(m_2)}{n^{m_2}(n+r+j-1)}-\frac{2(-1)^{n-1}}{n^{m_1+m_2}(n+r+j-1)}\right\}.\end{aligned}$$

Theorem 5.5 *For integers $m_1>0,m_2>0,m_3>0,k\geq 1,r\geq 0$, we have*

$$\begin{aligned}&(k-r)\sum_{n=1}^{\infty}\frac{\zeta_n(m_1)\,\zeta_n(m_2)\,\zeta_n(m_3)-\zeta_n(m_1+m_2+m_3)}{(n+r)(n+k)}\\&=\sum_{j=1}^{k-r}\sum_{n=1}^{\infty}\left\{\frac{\zeta_n(m_1)\,\zeta_n(m_2)}{n^{m_3}(n+r+j-1)}+\frac{\zeta_n(m_1)\,\zeta_n(m_3)}{n^{m_2}(n+r+j-1)}+\frac{\zeta_n(m_2)\,\zeta_n(m_3)}{n^{m_1}(n+r+j-1)}\right\}\\&\quad-\sum_{j=1}^{k-r}\sum_{n=1}^{\infty}\left\{\frac{\zeta_n(m_1)}{n^{m_2+m_2}(n+r+j-1)}+\frac{\zeta_n(m_2)}{n^{m_1+m_3}(n+r+j-1)}+\frac{\zeta_n(m_3)}{n^{m_1+m_2}(n+r+j-1)}\right\}.\end{aligned}$$

Some illustrative examples follow.

$$\sum_{n=1}^{\infty}\frac{\zeta_n(2)}{n(n+k)}=\frac{1}{k}\left\{\zeta(3)+\zeta(2)\,H_{k-1}-\sum_{i=1}^{k-1}\frac{H_i}{i^2}\right\},\tag{5.15}$$

$$\sum_{n=1}^{\infty}\frac{\zeta_n(3)}{n(n+k)}=\frac{1}{k}\left\{\zeta(4)+\zeta(3)\,H_{k-1}-\zeta(2)\,\zeta_{k-1}(2)+\sum_{i=1}^{k-1}\frac{H_i}{i^3}\right\},\tag{5.16}$$

$$\sum_{n=1}^{\infty}\frac{H_n^2}{n(n+k)}=\frac{1}{k}\left\{\begin{array}{l}3\zeta(3)+\dfrac{H_k^3+3H_k\zeta_k(2)+2\zeta_k(3)}{3}\\-\dfrac{H_k^2+\zeta_k(2)}{k}-\displaystyle\sum_{i=1}^{k-1}\frac{H_i}{i^2}+\zeta(2)\,H_{k-1}\end{array}\right\},\tag{5.17}$$

$$\sum_{n=1}^{\infty}\frac{H_n^3}{n\left(n+k\right)}=\frac{1}{k}\left\{\begin{array}{l}10\zeta\left(4\right)+\frac{1}{4}Y_4\left(k\right)-\frac{1}{k}Y_3\left(k\right)+4\zeta\left(3\right)H_{k-1}+\frac{1}{2}\zeta\left(2\right)\zeta_{k-1}\left(2\right)\\+\frac{3}{2}\zeta\left(2\right)H_{k-1}^2+\sum_{i=1}^{k-1}\frac{H_i}{i^3}-\frac{3}{2}\sum_{i=1}^{k-1}\frac{H_i^2+\zeta_i\left(2\right)}{i^2}-3\sum_{i=1}^{k-1}\frac{1}{i}\sum_{j=1}^{i}\frac{H_j}{j^2}\end{array}\right\}, \tag{5.18}$$

$$\begin{aligned}\sum_{n=1}^{\infty}\frac{H_n\zeta_n\left(2\right)}{n\left(n+k\right)}=&\frac{2\zeta\left(4\right)+2\zeta\left(3\right)H_{k-1}+\frac{1}{2}\zeta\left(2\right)H_{k-1}^2+\sum_{i=1}^{k-1}\frac{H_i}{i^3}}{k}\\&-\frac{\frac{1}{2}\zeta\left(2\right)\zeta_{k-1}\left(2\right)+\frac{1}{2}\sum_{i=1}^{k-1}\frac{H_i^2+\zeta_i\left(2\right)}{i^2}+\sum_{i=1}^{k-1}\frac{1}{i}\sum_{j=1}^{i}\frac{H_j}{j^2}}{k},\end{aligned} \tag{5.19}$$

$$\begin{aligned}\sum_{n=1}^{\infty}\frac{L_n^2\left(1\right)}{n\left(n+k\right)}=&\frac{3\zeta\left(2\right)\ln 2-\frac{9}{4}\zeta\left(3\right)+\ln^2 2\left(H_{k-1}+L_{k-1}\left(1\right)\right)-\zeta\left(2\right)L_{k-1}\left(1\right)}{k}\\&+\frac{\sum_{i=1}^{k-1}\frac{H_i}{i^2}+2\sum_{j=1}^{k-1}\frac{\left(-1\right)^{j-1}}{j}\sum_{i=1}^{j-1}\frac{H_i}{i}(-1)^{i-1}}{k},\end{aligned} \tag{5.20}$$

$$\begin{aligned}\sum_{n=1}^{\infty}\frac{H_nL_n\left(1\right)}{n\left(n+k\right)}=&\frac{\frac{3}{2}\zeta\left(2\right)\ln 2+\frac{1}{2}\zeta\left(2\right)H_{k-1}+\ln 2\left(\zeta_{k-1}\left(2\right)+L_{k-1}\left(2\right)\right)}{k}\\&-\frac{\frac{3}{8}\zeta\left(3\right)+\frac{1}{2}\ln^2 2L_{k-1}\left(1\right)+\sum_{i=1}^{k-1}\frac{L_i\left(1\right)}{i^2}(-1)^{i-1}}{k}\\&+\frac{2\ln 2\sum_{j=1}^{k-1}\frac{H_{j-1}+L_{j-1}\left(1\right)}{j}-\sum_{j=1}^{k-1}\frac{\left(-1\right)^{j-1}}{j}\sum_{i=1}^{j-1}\frac{L_i\left(1\right)}{i}}{k}\\&-\frac{1}{2k}\sum_{j=1}^{k-1}\frac{L_{j-1}^2\left(1\right)+\zeta_{j-1}\left(2\right)}{j}.\end{aligned} \tag{5.21}$$

Finally, we consider the following integral

$$\int_0^1 x^{m-1}\mathrm{Li}_s\left(x\right)\mathrm{Li}_t\left(x\right)\mathrm{Li}_r\left(x\right)dx\ \ \left(1\leq m,s,t,r\in\mathbb{Z}\right).$$

By using Cauchy product formula, it is easy to show that the left side equals

$$\mathrm{Li}_s\left(x\right)\mathrm{Li}_t\left(x\right)=\sum_{n=1}^{\infty}\sum_{k=1}^{n}\frac{x^{n+1}}{k^s(n-k+1)^t}. \tag{5.22}$$

Applying the following formula

$$\frac{1}{x^s(1-x)^t}=\sum_{j=1}^{s}\frac{A_j^{(s,t)}}{x^j}+\sum_{j=1}^{t}\frac{B_j^{(s,t)}}{(1-x)^j},\quad s,t\geq 0,s+t\geq 1, \tag{5.23}$$

to (5.22), we have

$$\mathrm{Li}_s(x)\,\mathrm{Li}_t(x)=\sum_{j=1}^{s}A_j^{(s,t)}\sum_{n=1}^{\infty}\frac{\zeta_n(j)}{n^{s+t-j}}x^n+\sum_{j=1}^{t}B_j^{(s,t)}\sum_{n=1}^{\infty}\frac{\zeta_n(j)}{n^{s+t-j}}x^n$$
$$-\left(\sum_{j=1}^{s}A_j^{(s,t)}+\sum_{j=1}^{t}B_j^{(s,t)}\right)\mathrm{Li}_{s+t}(x),\tag{5.24}$$

where $A_j^{(s,t)}=\binom{s+t-j-1}{s-j},B_j^{(s,t)}=\binom{s+t-j-1}{t-j}$. Multiplying (5.24) by x^{m-1} and integrating over (0,1), we can get

$$\begin{aligned}&\int_0^1 x^{m-1}\mathrm{Li}_s(x)\,\mathrm{Li}_t(x)dx\\&=\sum_{j=1}^{s}A_j^{(s,t)}\sum_{n=1}^{\infty}\frac{\zeta_n(j)}{n^{s+t-j}(n+m)}+\sum_{j=1}^{t}B_j^{(s,t)}\sum_{n=1}^{\infty}\frac{\zeta_n(j)}{n^{s+t-j}(n+m)}\\&\quad-\left(\sum_{j=1}^{s}A_j^{(s,t)}+\sum_{j=1}^{t}B_j^{(s,t)}\right)\left\{\sum_{i=1}^{s+t-1}\frac{(-1)^{i-1}}{m^i}\zeta(s+t+1-i)+\frac{(-1)^{s+t-1}}{m^{s+t}}H_m\right\}.\end{aligned}\tag{5.25}$$

Multiplying (5.24) by $x^{m-1}\mathrm{Li}_r(x)$ and integrating over (0,1), we can obtain

$$\begin{aligned}&\int_0^1 x^{m-1}\mathrm{Li}_s(x)\,\mathrm{Li}_t(x)\,\mathrm{Li}_r(x)dx\\&=\sum_{j=1}^{s}A_j^{(s,t)}\sum_{n=1}^{\infty}\frac{\zeta_n(j)}{n^{s+t-j}}\left\{\sum_{i=1}^{r-1}\frac{(-1)^{i-1}}{(n+m)^i}\zeta(r+1-i)+\frac{(-1)^{r-1}}{(n+m)^r}H_{n+m}\right\}\\&\quad+\sum_{j=1}^{t}B_j^{(s,t)}\sum_{n=1}^{\infty}\frac{\zeta_n(j)}{n^{s+t-j}}\left\{\sum_{i=1}^{r-1}\frac{(-1)^{i-1}}{(n+m)^i}\zeta(r+1-i)+\frac{(-1)^{r-1}}{(n+m)^r}H_{n+m}\right\}\\&\quad-\left(\sum_{j=1}^{s}A_j^{(s,t)}+\sum_{j=1}^{t}B_j^{(s,t)}\right)\int_0^1 x^{m-1}\mathrm{Li}_{s+t}(x)\,\mathrm{Li}_r(x)dx.\end{aligned}\tag{5.26}$$

In the evaluation of $\int_0^1 x^{m-1}\mathrm{Li}_s(x)\,\mathrm{Li}_t(x)\,\mathrm{Li}_r(x)dx$ we require the closed form representation of

$$\sum_{n=1}^{\infty}\frac{H_{n+m}\zeta_n(r)}{n^s(n+m)^t}.$$

From (5.23), we know that

$$\frac{1}{(n+r)^s(n+k)^t}=(-1)^s\sum_{l=1}^{s}(-1)^l\frac{A_l^{(s,t)}}{(n+r)^l(k-r)^{s+t-l}}+(-1)^s\sum_{l=1}^{t}\frac{B_l^{(s,t)}}{(n+k)^l(k-r)^{s+t-l}},\tag{5.27}$$

where $0 \leq r < k \in \mathbb{Z}$. Now, we evaluate the following Euler type sums

$$\sum_{n=1}^{\infty} \frac{\zeta_{n+m_1}(p_1)\,\zeta_{n+m_2}(p_2)}{(n+r)(n+k)},\ 1 < p_1, p_2 \in \mathbb{Z},\ 0 \leq r < k \in \mathbb{Z},$$

with $0 \leq m_1, m_2 \in \mathbb{Z}$. Considering the power series function

$$y = \sum_{n=1}^{\infty} \zeta_{n+m_1}(p_1)\,\zeta_{n+m_2}(p_2) x^{n-1}, x \in (-1,1).$$

By the definition of $\zeta_n(r) = \sum_{j=1}^{n} \frac{1}{j^r}$, we deduce that

$$\begin{aligned}
&(1-x)\sum_{n=1}^{\infty} \zeta_{n+m_1}(p_1)\,\zeta_{n+m_2}(p_2) x^{n-1} \\
&= \zeta_{m_1}(p_1)\,\zeta_{m_2}(p_2) + \sum_{n=1}^{\infty} \left\{ \frac{\zeta_{n+m_1}(p_1)}{(n+m_2)^{p_2}} + \frac{\zeta_{n+m_2}(p_2)}{(n+m_1)^{p_1}} - \frac{1}{(n+m_1)^{p_1}(n+m_2)^{p_2}} \right\} x^{n-1}. \quad (5.28)
\end{aligned}$$

Multiplying (5.28) by $\frac{x^r - x^k}{1-x}$ and integrating over (0,1), we can obtain

$$\begin{aligned}
&(k-r)\sum_{n=1}^{\infty} \frac{\zeta_{n+m_1}(p_1)\,\zeta_{n+m_2}(p_2)}{(n+r)(n+k)} \\
&= \zeta_{m_1}(p_1)\,\zeta_{m_2}(p_2)\,(H_k - H_r) \\
&\quad + \sum_{i=1}^{k-r}\sum_{n=1}^{\infty} \left\{ \frac{\zeta_{n+m_1}(p_1)}{(n+m_2)^{p_2}} + \frac{\zeta_{n+m_2}(p_2)}{(n+m_1)^{p_1}} - \frac{1}{(n+m_1)^{p_1}(n+m_2)^{p_2}} \right\} \frac{1}{n+r+i-1}. \quad (5.29)
\end{aligned}$$

Using (5.6),(5.27) and noting that

$$\begin{aligned}
\sum_{n=1}^{\infty} \frac{\zeta_{n+m}(p)}{(n+r)(n+k)} &= \sum_{n=1}^{\infty} \frac{\zeta_n(p) + \sum_{i=1}^{m} \frac{1}{(n+i)^p}}{(n+r)(n+k)} \\
&= \sum_{n=1}^{\infty} \frac{\zeta_n(p)}{(n+r)(n+k)} + \sum_{i=1}^{m}\sum_{n=1}^{\infty} \frac{1}{(n+i)^p (n+r)(n+k)} \\
&= \sum_{n=1}^{\infty} \frac{\zeta_n(p)}{(n+r)(n+k)} + \sum_{i=1}^{m} \frac{1}{k-r} \sum_{n=1}^{\infty} \left\{ \frac{1}{(n+i)^p (n+r)} - \frac{1}{(n+i)^p (n+k)} \right\}, \quad (5.30)
\end{aligned}$$

we can obtain some closed form representation of $\sum_{n=1}^{\infty} \frac{\zeta_{n+m_1}(p_1)\,\zeta_{n+m_2}(p_2)}{(n+r)(n+k)}$.

Acknowledgments. The authors would like to thank the anonymous referee for his/her helpful comments, which improve the presentation of the paper.

References

[1] H. Alzer, D. Karayannakis, H.M. Srivastava. *Series representations for some mathematical constants*. J. Math. Anal. Appl., 2006, **320**: 145–162.

[2] George E. Andrews, Richard Askey, Ranjan Roy. *Special Functions*. Cambridge University Press., 2000: 481-532.

[3] David H. Bailey, Jonathan M. Borwein and Roland Girgensohn. *Experimental evaluation of Euler sums*. Experimental Mathematics., 1994, **3**(1): 17-30.

[4] David H. Bailey, Jonathan M. Borwein, Richard E. Crandall. *Computation and theory of extended Mordell-Tornheim-Witten sums*. Math. Comp., 2014, **83**(288): 1795-1821.

[5] B. C. Berndt. *Ramanujans Notebooks, Part I*. Springer-Verlag, New York., 1985.

[6] B. C. Berndt. *Ramanujans Notebooks, Part II*. Springer-Verlag, New York., 1989.

[7] J. Blumlein, S. Kurth. *Harmonic sums and Mellin transforms up to two loop order*. Phys. Rev. D., 1999, **60**: 014018.

[8] David Borwein, Jonathan M. Borwein and Roland Girgensohn. *Explicit evaluation of Euler sums*. Proc. Edinburgh Math., 1995, **38**: 277-294.

[9] J. Borwein, P. Borwein, R. Girgensohn, S. Parnes. *Making sense of experimental mathematics*. Mathematical Intelligencer., 1996, **18**(4): 12-18.

[10] Jonathan M. Borwein, David M. Bradley, David J. Broadhurst, Petr. Lisonk. *Special values of multiple polylogarithms. Trans*. Amer. Math. Soc., 2001, **353**(3): 907-941.

[11] J. M. Borwein, I. J. Zucker, J. Boersma. *The evaluation of character Euler double sums*. Ramanujan J., 2008, **15** (3): 377-405.

[12] J. M. Borwein, R. Girgensohn, *Evaluation of triple Euler sums*, Electron. J. Combin., 1996: 2-7.

[13] M. W. Coffey, *On some log-cosine integrals related to* $\zeta(3), \zeta(4)$, *and* $\zeta(6)$, J. Comput. Appl. Math., 2003, **159**: 205–215.

[14] M. W. Coffey, *On one-dimensional digamma and polygamma series related to the evaluation of Feynman diagrams*, J. Comput. Appl. Math., 2005, **183**: 84–100.

[15] M. W. Coffey, *On a three-dimensional symmetric Ising tetrahedron and contributions to the theory of the dilogarithm and Clausen functions*, Journal of Mathematical Physics., 2008, **49**(4): 542–555.

[16] M. W. Coffey, Lubbers N. *On generalized harmonic number sums*. Applied Mathematics and Computation., 2010, **217**(2): 689-698.

[17] Ayhan Dil, Khristo N. Boyadzhiev, *Euler sums of hyperharmonic numbers*, J. Number Theory., 2015, **147**: 490-498.

[18] Minking Eie, Chuan-Sheng Wei. *Evaluations of some quadruple Euler sums of even weight*. Functions et Approximatio., 2012, **46**(1): 63-67.

[19] Philippe Flajolet, Bruno Salvy. *Euler sums and contour integral representations*. Experimental Mathematics., 1998, **7**(1): 15–35.

[20] Pedro freitas. *Integrals of polylogarithmic functions, recurrence relations, and associated Euler sums*. Mathematics of Computation,. 2005, **74**(251): 1425-1440.

[21] Comtet L. Advanced combinatorics, Boston: D Reidel Publishing Company, 1974.

[22] C. Markett. *Triple Sums and the Riemann Zeta Function*. J. Number Theory, 1994, **48**(2): 113-132.

[23] I. Mezö, A. Dil. *Hyperharmonic series involving Hurwitz zeta function*. Journal of Number Theory, 2010, **130**(2): 360–369.

[24] I. Mezö. *Nonlinear Euler sums*. Pacific J. Math., 2014, **272**: 201-226.

[25] T.M. Rassias, H.M. Srivastava. *Some classes of infinite series associated with the Riemann zeta function and polygamma functions and generalized harmonic numbers*. Appl. Math. Comput., 2002, **131**: 593–605.

[26] A. Sofo. *Integral forms of sums associated with harmonic numbers*. Applied Mathematics and Computation., 2009, **207**(2): 365-372.

[27] A. Sofo. *Quadratic alternating harmonic number sums*. J. Number Theory., 2015, **154**: 144-159.

[28] A. Sofo. *Integral identities for sums*. Math. Commun., 2008, **13**: 303-309.

[29] A. Sofo, H.M. Srivastava. *Identities for the harmonic numbers and binomial coefficients*. Ramanujan J., 2011, **25**: 93-113.

[30] A. Sofo. *Harmonic sums and integral representations*. J. Appl. Anal., 2010, **16**: 265-277.

[31] A. Sofo. *Harmonic number sums in closed form*. Math. Commun., 2011, **16**: 335-345.

[32] A. Sofo. *Shifted harmonic sums of order two*. Commun. Korean Math. Soc., 2014, **29**(2): 239-255.

[33] Ce Xu, Jinfa Cheng. *Some Results On Euler Sums*. Functions et Approximatio., 2016, **54**(1): 25-37.

[34] Ce Xu, Yuhuan Yan, Zhijuan Shi. *Euler sums and integrals of polylogarithm functions*. J. Number Theory., 2016, **165**: 84-108.

[35] Ce Xu, Yingyue Yang, Jianwen Zhang. *Explicit evaluation of quadratic Euler sums*. Int. J. Number Theory., 2016, in press.

[36] D. Zagier.*Values of zeta functions and their applications*. First European Congress of Mathematics, Volume II, Birkhauser, Boston., 1994, (120): 497-512.

[37] D. Zagier. *Evaluation of the multiple zeta values* $\zeta(2,...,2,3,2,...,2)$. Annals of Mathematics., 2012, **2**(2): 977-1000.